



PARALLEL SESSIONS: ORAL COMMUNICATIONS

3rd World Congress of Health Research

Viseu - Portugal, 29-30 September 2016

1. ADULT HEALTH

CHARACTERIZATION OF PATIENTS UNDERGOING NONINVASIVE VENTILATION ADMITTED IN UNIT INTERMEDIATE CARE

Matilde Martins^a, Patrícia Ribas^b, Joana Sousa^c, Andreia Cunha^c, Norberto Silva^d, Teresa Correia^a

^aHealth School of Polytechnic Institute of Bragança, Sports Sciences, Health Sciences and Human Development, Portugal;

^bClinic of Arrifana, Portugal; ^cHealth School of Polytechnic Institute of Bragança, Portugal; ^dHealth School of Polytechnic Institute of Bragança, Unit of the Northeast Local Health Bragança, Portugal.

Contact details: matildemartins@ipb.pt

Introduction: Non-invasive ventilation (NIV) is the application of a ventilatory support without resorting to invasive methods. Today it's considered a credible therapeutic option, with enough scientific evidence to support its application in various situations and clinical settings related to the treatment of acute respiratory disease, as well as chronic respiratory disease.

Objectives: Characterize patients undergoing NIV admitted in Unit Intermediate Care (ICU) in the period from October 1st 2015 to June 30th 2016.

Methods: Prospective study conducted in ICU between October 2015 and June 2016. In this study were included all patients hospitalized in this unit (ICU) and in that time period a sample of 57 participants was obtained. As data collection instruments we used a questionnaire for sociodemographic and clinical data and the Braden scale.

Results: Participants were mostly male 38 (66.7%), the average age 69.5 ± 11.3 years, ranging between 43 and 92 years. They weighed on average 76.6 kg (52 and 150), with an average body mass index of 28.5 kg/m^2 (20 to 58.5). With skin intact 28 (49.1%) with abnormal perfusion 12 (21.1%), with altered sensitivity 11 (19.3%) and a high risk of ulcer on the scale of Braden 37 (65%). The admission diagnosis was respiratory failure 33 (57.3%) and had different backgrounds. We used reused mask 53 (93.0%), the average time of NIV was 7.1 days (1-28), 4.8 days of hospitalization (1-18) and an average of 7.8 IPAP pressure. 11 (19.3%) of the participants developed face ulcer pressure.

Conclusions: The NIV is used in patients with advanced age, obesity, respiratory failure and high risk of face ulcer development.

Keywords: Patients. Noninvasive ventilation.

TELEPHONE SUPPORT LINE OF RHEUMATOLOGY DEPARTMENT: A 4.5 YEARS EXPERIENCE

Eduardo Santos^a, Andréa Marques^b

^aAssociate Research at the Portugal Centre for Evidence-Based Practice (PCEBP): a Collaborating Centre of the Joanna Briggs Institute - Health Sciences Research Unit: Nursing (UICISA: E), Nursing School of Coimbra (ESEnfC); ^bRheumatology department, Centro Hospitalar e Universitário de Coimbra, EPE.

Contact details: ejf.santos87@gmail.com

Introduction: In many health services in developed countries, the telephone support has been used as an innovative approach to providing care and answering questions by nurses, developing especially in specific areas such as rheumatology (Brown et al., 2006).

Objectives: To analyze the profile of users and the main reasons of patients that uses the Telephone Support Line of Day Hospital and early arthritis consultation of the CHUC Rheumatology Department, EPE.

Methods: This is a retrospective descriptive-correlational study with a convenience sample of 448 calls. For continuous variables we used the t-student test, for dichotomous variables the χ^2 test and, finally, we performed a cluster analysis by the TwoStep Cluster method through the log-likelihood distance.

Results: Users have an average age of 44.8 years being predominantly male (58%). The cluster analysis allowed to create three groups whose profile shows that the cluster 1 ($n = 96/21.4\%$) are formed by "older" adults that need to clarify mainly doubts about the results of auxiliary examinations tests (AET)/prescriptions and side effects/complications; the cluster 2 ($n = 232/51.8\%$) are formed by young adults that call to change appointments and mainly to clarify doubts on the results of AET/prescriptions; finally, the cluster 3 ($n = 120/26.8\%$) are formed by "middle age" adults that want to clarify doubts about medication and disease management.

Conclusions: Incoming calls cover a wide range of ages and all kinds of rheumatic diseases. This study allows us to identify in which groups it is necessary to make a more detailed educational

ministered questionnaire validated for adolescents was conducted by Di Pietro (2002). The sample is composed of 44% male and 56% female adolescents, with an age mean of 15.54. 61.2% of the boys and 83.6% of the girls stated to be concerned with their weight. The main reasons given for this concern were health associated with aesthetics (29.1% of boys and 38.5% of girls). The results show that 12.8% of boys and 23.5% of girls are dissatisfied with their body image. The study revealed that the variable gender is statistically moderating in relation to the variables: body image perception and concern with weight: female adolescents show a higher dissatisfaction with their body image (0.003) and a bigger concern with weight (0.000). The results point out towards the need for assessment/intervention in this population as body image represents a paramount issue in adolescence, with the body image self-perception being strongly associated with the biopsychosocial maturing process, which interferes with their level of health and personal and social development.

Keywords: Body image. Adolescents. Youth health.

PARENTAL FEELINGS FACE TO HOSPITALIZATION OF THE CHILD

Celeste Antão, Nicole Rodrigues, Filomena Sousa

Polytechnic Institute of Bragança, Bragança, Portugal.

Contact details: celeste@ipb.pt

Introduction: The hospitalization of a child carries with it major changes, either in the child's life or parent. Illness and hospitalization constitute a crisis situation for both the child and the family as it incorporates financial, psychological, relational and social changes. The purpose of this work and analyze the satisfaction and parents' perception on the importance of nursing care during hospitalization and feelings experienced by parents.

Objectives: To know the opinions and feelings of the parents in relation to the admission of the child in the service of Pediatrics at North of Portugal.

Methods: Integrated this study parents of children hospitalized in the pediatric inpatient at the Hospital Trás-os-Montes e Alto Douro, Vila Real, whose children were in the service to at least more than 48 hours, In total 33 escorts that replied to the questionnaire. For such a study we chose the type quantitative descriptive. We proceeded to collect data by conducting a questionnaire, and the results of it were drafted in the SPSS program through Quantitative model with exploratory qualitative approach.

Results: The age of the participants is between the 21 and 46 years - old, 75.8% of respondents were married, 78.8% of children was hospitalized in the first time. We conclude that the mother is who else accompanies the child in the hospital. We had positive feedback regarding the interaction parent-nurses since many parents stressed the good relationship with the nursing team (It reported that "Nurses" and "Nurses and auxiliaries" were who else helped them, with 30.3% and 21.2% respectively). The feelings more mentioned were: anxiety -21 answers, sadness-20, fear-15 and hope with 15. The conditions offered to the parents were considered good for 60.6%.

Keywords: Nursing. Parenting. Feelings. Hospitalization of children.

HEARING LOSS AND CHILDREN'S BALANCE

Maria Inês Araújo, Lucie Esteves

Coimbra Health School, Portugal.

Contact details: ines@estescoimbra.pt

Introduction: Body balance is essential for normal development and most of the child's daily activities. For this to happen smoothly,

there must be consistency between the three sensory systems: visual, proprioception and vestibular.

Objectives: To assess the body balance and vestibular system in children with severe to profound hearing, with and without hearing impairment and children without hearing loss.

Methods: For the collection of information Posturography Computerised (PC), Clinical Test of Sensory Integration and Balance (mCTSIB) and the one-leg support test were applied. It was held further Fukuda/Unterberger test, Subjective Visual Vertical (SVV) and applied research of spontaneous and decentered nystagmus. For this study, we used a sample constituted by 32 children, 18 with hearing disorders and 15 with normal hearing, aged 12 to 17 years.

Results: We found that no significant differences between children with hearing disorders and normal hearing in any of the tests. The results revealed no significant differences between children with hearing disorders and normal hearing in any of the tests. Although, when compared the children who practice sports twice or three times per week in both groups, it is found statistically relevant variations on mCTSIB test.

Conclusions: In this study children with severe to profound hearing loss started walking with the age to normal hearing children, do not have higher body instability. However, regular sports practice causes a higher body balance in normal hearing children compared with children with hearing loss.

Keywords: Children. Hearing loss. Body balance.

ADOLESCENTS WITH CYSTIC FIBROSIS AND THEIR INFORMATION DESK: INTERNET, PARENTS OR HEALTH PROFESSIONALS?

Conceição Reisinho, Bárbara Gomes

Escola Superior de Enfermagem do Porto, Portugal.

Contact details: creisinho@esenf.pt

Introduction: The development of diagnostic and treatment options for cystic fibrosis has had a positive impact on children's survival rates but also led to the appearance of more disease-related problems. The children's response mechanisms and ability to adapt to this new situation are consistent with the personal physical, psychological, emotional and social development and to society status. In the adolescents with Cystic Fibrosis (CF), besides the age related developmental changes, several physical and psychosocial needs are identified, according to the disease characteristics.

Objectives: To identify the sources of information on cystic fibrosis searched for adolescents.

Methods: A qualitative study was performed, using semi-structured interviews for data collection. The sample comprised 16 adolescents with CF diagnosed for more than one year. Adolescents were aged between 11 and 23 years, living in the northern region of Portugal.

Results: The internet was the main tool used to search for disease-related information, in particular on symptoms, prognosis and treatments. Adolescents accessed information using a general web search tool, rather than a specific web site. Parents and health professionals, especially the practitioner, who accompany the adolescent since diagnosis, were also considered a very reliable source of information.

Conclusions: This study results showed that adolescents are interested on information about their health. Participants also refer to graphically sensitive images and information contents on prognosis and treatment options, likely to undermine the adherence to treatment. Nurses are professionals able to establish a closer connection with this population, thus their interventions should focus on providing guidance and relevant information.

Keywords: Adolescents. Cystic fibrosis. Information.