



## VARIATION OF LIPIDS INDEXES IN POLLEN WITH ITS BOTANICAL ORIGIN

O Anjos<sup>1, 2</sup>, G Campos Maria<sup>3, 4</sup>, T Dias<sup>5</sup>, LM Estevinho<sup>5, 6</sup>

PMIO 2017; 4(S 01): S1-S20. DOI: 10.1055/s-0037-1608202

<sup>1</sup>Instituto Politécnico de Castelo Branco, Castelo Branco, Portugal

<sup>2</sup>Centro de Biotecnologia de Plantas da Beira Interior, Castelo Branco, Portugal

<sup>3</sup>Observatory of Herb-Drug Interactions/Faculty of Pharmacy, University of Coimbra, Health Sciences Campus, Coimbra, Portugal

<sup>4</sup>Coimbra Chemistry Centre (CQC, FCT Unit 313) (FCTUC) University of Coimbra, Coimbra, Portugal

<sup>5</sup>Agricultural College of Bragança Polytechnic Institut of Bragança, Bragança, Portugal

<sup>6</sup>Center of Molecular and Environmental Biology, University of Minho, Braga, Portugal

Bee pollen is one of nature's healthful food products with promising nutritional and therapeutic properties due to its chemical composition, particularly its protein content, which includes almost all the essential amino acids. Nevertheless the composition in lipids it is not completely known and supposedly will be highly variable depending on the floral origin. As such, this parameter must be ascertained in line with the research for bioactivity [1].


The aim of this work was to evaluate some health-related lipid indexes of bee pollen, namely omega-6/omega-3 fatty acid ratio (n-6/n-3); polyunsaturated fatty acid/saturated fatty acid ratio (PUFA/SAT), atherogenic index (AI) and thrombogenic index (TI) in samples harvested in Portugal. The selected parameters were calculated from the Fatty-Acid Profile, which was determined as previously reported by Bárbara et al. [3].

Bee pollen samples, after harvest, were cleaned and frozen at -20 °C and were codified according the predominant pollen. Figure 1 associates the lipid indexes of the different samples with their botanical origins.

n-6/n-3 and PUFA/SAT ratios were in within the limits recommended by World Health Organization (below 4.0 and above 0.45, respectively), suggesting that bee pollen is a good product with the nutritional point of view, with potential beneficial effects for the consumer's health. Also, both AI and TI indexes of this natural product were low, even though this effect depended on bee pollen's botanical origin (Figure 1).

[1] Campos MG, Olena L. and Anjos O. 2016. Chapter 3, Chemical Composition of Bee Pollen. In Cardoso SM & Silva AMS, Chemistry, biology and potential applications of honeybee plant-derived products, Bentham Science Publishers, United Arab. Pp. 67 – 88 (22).

[2] Bárbara, M.S, Machado, C.S, Sodr , G.D.S, Dias, L.G, Estevinho, L.M, & de Carvalho, C.A.L. Molecules, 2015,20, 12525 – 12544.



**Figure 1** – Principal component analysis for the bee pollen sample representing the relation between samples and lipid indexes: Ru (sample with 90% of the pollen grains from *Rubus* spp.); CA (sample with 91% of *Castanea sativa*); Ci (sample with 90% of *Cistus* spp.); Le (sample with 99% of *Leontodon* spp.); Cv (sample with 90% of *Cytisus* spp.); Ec (sample with 89% of *Echium* spp.); and Er (sample with 90% of *Erica* spp.).