

SELECTED, EXTENDED AND REVISED CONTRIBUTIONS FROM THE INTERNATIONAL SYMPOSIUM OCCUPATIONAL SAFETY AND HYGIENE, GUIMARÃES, PORTUGAL, 23–24 MARCH 2016

Occupational Safety and Hygiene IV

Editors

Pedro M. Arezes

University of Minho, Guimarães, Portugal

João Santos Baptista

University of Porto, Porto, Portugal

Mónica P. Barroso, Paula Carneiro, Patricio Cordeiro & Nélson Costa

University of Minho, Guimarães, Portugal

Rui B. Melo

University of Lisbon, Lisbon, Portugal

A. Sérgio Miguel

University of Minho, Guimarães, Portugal

Gonçalo Perestrelo

SPOSHO, Porto, Portugal



CRC Press

Taylor & Francis Group

Boca Raton London New York Leiden

CRC Press is an imprint of the
Taylor & Francis Group, an **informa** business

A BALKEMA BOOK

CRC Press/Balkema is an imprint of the Taylor & Francis Group, an informa business

© 2016 Taylor & Francis Group, London, UK

Typeset by V Publishing Solutions Pvt Ltd., Chennai, India

Printed and bound in Great Britain by CPI Group (UK) Ltd, Croydon, CR0 4YY

All rights reserved. No part of this publication or the information contained herein may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, by photocopying, recording or otherwise, without written prior permission from the publisher.

Although all care is taken to ensure integrity and the quality of this publication and the information herein, no responsibility is assumed by the publishers nor the author for any damage to the property or persons as a result of operation or use of this publication and/or the information contained herein.

Published by: CRC Press/Balkema

P.O. Box 11320, 2301 EH Leiden, The Netherlands

e-mail: Pub.NL@taylorandfrancis.com

www.crcpress.com – www.taylorandfrancis.com

ISBN: 978-1-138-02942-2 (Hbk)

ISBN: 978-1-315-62896-7 (eBook PDF)

Table of contents

Foreword	xi
Occupational exposure assessment to particles with task-based approach <i>S. Viegas, M. Almeida-Silva, T. Faria, M. Dos Santos & C. Viegas</i>	1
Trends of Volatile Organic Compounds in different indoor microenvironments: A review <i>I. Paciência, J. Madureira, J. Cavaleiro Rufo, E. Oliveira Fernandes, A. Moreira & J.P. Teixeira</i>	7
Climate changes influence in occupational exposure to fungi and mycotoxins <i>C. Viegas, M. Meneses & S. Viegas</i>	11
Changes of air temperature and its effects on student's health <i>J.C.F. Siqueira, L.B. da Silva, A.S. Coutinho & R.M. Rodrigues</i>	17
Biomechanical analysis of the manual handling of fertilizers in silvicultural operations, in mountainous regions <i>L.J. Minette, S. Schettino, A.P. de Souza, F.L. da Cunha Marzano & A.C.M. Camarinha</i>	21
Exposure to emissions from cleaning products in primary schools: A test chamber study <i>J. Cavaleiro Rufo, J. Madureira, I. Paciência, J.R.B. Sousa, E. Oliveira Fernandes, K. Slezakova, M.C. Pereira, L. Aguiar, J.P. Teixeira, M. Pinto, L. Delgado & A. Moreira</i>	25
Analysis of overload in the musculoskeletal system of women developing repetitive tasks in fluid filling process in chemical industry <i>L. Monteiro, J. Santos, V. Santos, V. Franca & O. Alsina</i>	29
Communication strategies in Occupational Safety and Health training for foreign university students: A case study <i>B. Mrugalska & T. Ahram</i>	35
The Portuguese anthropometric database comparison <i>A.L. Cabral</i>	41
Human systems engineering approach for safety and risk management of complex systems design <i>T. Ahram, W. Karwowski & C. Falcão</i>	45
Assessment of gaseous pollutants in a preschool environment <i>M. Oliveira, K. Slezakova, M.C. Pereira, C. Delerue-Matos & S. Morais</i>	51
Influence of psychosocial factors in the pain symptoms <i>J.M.N. Silva, W.K.S. Leite, R.E.C. Lucas, J.G.B. Fernandes, C.A. Falcão, E.L. Souza, L.B. Silva & G. Colaço</i>	57
Development of ergonomic sofas for labor rest areas <i>S. Schettino, R. Hourani, L.J. Minette & A.C.M. Camarinha</i>	63
Performance evaluation in teaching environments on climate change <i>R. Rodrigues, L. Silva, E. Souza, A. Souza & F. Brito</i>	69
Firefighter's occupational exposure to PM _{2.5} and Polycyclic Aromatic Hydrocarbons <i>M. Oliveira, K. Slezakova, M.C. Pereira, A. Fernandes, J.A. Vaz, C. Delerue-Matos & S. Morais</i>	73
Influence of temperatures' variation on students' productivity <i>F. Brito, R. Rodrigues, L. Silva, E. Souza, R. Souza & A. Coutinho</i>	77

Firefighter's occupational exposure to PM_{2.5} and Polycyclic Aromatic Hydrocarbons

M. Oliveira

REQUIMTE-LAQV, Instituto Superior de Engenharia do Porto, Porto, Portugal
LEPABE, Faculdade de Engenharia, Universidade do Porto, Porto, Portugal

K. Slezakova & M.C. Pereira

LEPABE, Faculdade de Engenharia, Universidade do Porto, Porto, Portugal

A. Fernandes & J.A. Vaz

Escola Superior de Saúde, Instituto Politécnico de Bragança, Bragança, Portugal

C. Delerue-Matos & S. Morais

REQUIMTE-LAQV, Instituto Superior de Engenharia do Porto, Porto, Portugal

ABSTRACT: This study collected the personal PM_{2.5} air fraction in fifteen healthy and no-smoking firefighters during their normal shift inside four Portuguese fire stations. Indoor PM_{2.5} levels varied between 0.05 to 1.04 µg/m³. Polycyclic Aromatic Hydrocarbons (PAHs) are known for their ubiquity and toxicity, being some of them classified as carcinogenic and possible carcinogens to humans. Firefighters' personal PM_{2.5}-bound total PAH concentrations ranged between 35.8 to 294 ng/m³ with total carcinogenic PAHs accounting with 12% to the total PAHs. Benzo[a]pyrene, the PAH biomarker of carcinogenicity, was detected in levels ranging from 6.74 × 10⁻² to 1.00 ng/m³.

1 INTRODUCTION

Firefighting is one of the most hazardous yet the least studied occupations in terms of exposures and their relationship to occupational diseases. Firefighters are chronically exposed to complex mixtures of particles, smoke and products of incomplete combustion which contain a high number of substances such as carbon monoxide and dioxide, nitrogen oxides, respirable Particulate Matter (PM), carbonyls (methanol, formaldehyde, 2,3-butanedione, acetaldehyde, acetone, methylethylketone), volatile (benzene, toluene, acetonitrile), and semi-volatile organic compounds including polychlorinated biphenyls, polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, Polycyclic Aromatic Hydrocarbons (PAHs), nitro-PAHs, and nitro-PAHs lactones (Estrellan & Lino 2010; Lemieux 2004; Lewtas 2007; Reisen et al. 2006).

Particulate matter is a mixture of solid or solid/liquid particles suspended in air that vary in size, shape, origin, and chemical composition (WHO 2006). PM with aerodynamic diameter lower than 2.5 µm, PM_{2.5}, is among the most relevant size fractions because it can reach lower parts of res-

piratory tract where it deposits in the conducting air ways and gas exchange regions of lungs. PM is originated from a variety of outdoor natural (pollen, spores, bacteria, plant, animal debris, and suspended crustal materials) and anthropogenic (combustion of fossil fuels, biomass burning, and ammonia emissions from agricultural operations) sources (WHO 2006). Indoor sources of PM include human activities such as cleaning, cooking, and combustion processes such as wood and fossil fuel burning, or tobacco smoke.

PAHs are one of the most relevant pollutants that are found adsorbed to PM surface. PAHs are classified as persistent organic pollutants (WHO 2013) with sixteen of them being regarded as priority pollutants by the USA Environmental Protection Agency (USEPA 2005). Among the 16 USEPA priority PAHs, benzo[a]pyrene is the only compound classified as carcinogenic to humans (i.e. group 1) by the International Agency for Research on Cancer. Naphthalene, benz[a]anthracene, benzo[b]fluoranthene, benzo[j]fluoranthene, benzo[k]fluoranthene, chrysene, and indeno[1,2,3-c,d]pyrene are classified as possibly carcinogenic to humans (group 2B; IARC 2002, 2010). Additionally, some studies have also included dibenzo[a,l]

pyrene and dibenz[a,h]anthracene because they are regarded as probable carcinogens to humans due to their higher carcinogenic potency than benzo[a]pyrene (Okona-Mensah et al. 2005; Oliveira et al. 2015). The presence of PAHs in the environment is primarily due to emissions from incomplete combustion of carbon fuels from natural, industrial, commercial, vehicular, and residential sources. When they are not actively suppressing fires on-duty firefighters stay, live and work at their fire stations. Scarce information is available regarding firefighters occupational exposure at fire stations.

The present work aims to assess firefighters' personal exposure to indoor $PM_{2.5}$ and $PM_{2.5}$ -bound PAHs at four Portuguese fire stations.

2 MATERIAL AND METHODS

2.1 Sampling

Fifteen healthy and no-smoking firefighters were asked to wear a personal air pump during four consecutive hours during their regular work shift indoors at four fire stations located in Bragança, north of Portugal. $PM_{2.5}$ -bound PAHs were collected on polytetrafluoroethylene membrane filters with polymethylpentene support ring (2 μ m porosity, SKC Ltd., United Kingdom) by constant flow sampler. $PM_{2.5}$ masses were determined gravimetrically according to Slezakova et al. (2014). Briefly, the initial mean mass of the blank filter was subtracted from the final mean mass of the exposed filter; the difference was then divided by the total volume of air that passed through the filter (at 25°C and 101.3 kPa). After the sampling, filters were stored in a freezer (-20°C) before consequent chemical analysis.

2.2 PAHs extraction and quantification

Sixteen USEPA priority PAHs, namely naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene anthracene, fluoranthene, pyrene, benz[a]anthracene, chrysene, benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[a]pyrene, dibenz[a,h]anthracene, benzo[g,h,i]perylene, and indeno[1,2,3-cd]pyrene, as well as dibenzo[a,l]pyrene and benzo[j]fluoranthene (recommended by EU Directive 2004/107/EC) were extracted from $PM_{2.5}$ by microwave assisted extraction according to the previously validated methodology (Castro et al. 2009, 2011). Extracts were analyzed using a Shimadzu LC system (Shimadzu Corporation, Kyoto, Japan) equipped with a LC-20 AD pump, DGU-20 AS degasser and photodiode array SPD-M20 A (PAD) and fluorescence RF-10 AXL (FLD) detectors on line. Separation of the compounds was performed in a C18 column (CC 150/4 Nucleosil

100-5 C18 PAH, 150 \times 4.0 mm; 5 μ m particle size; Macherey-Nagel, Duren, Germany) maintained at room temperature (20 \pm 1°C). The total run time was 40 min with a flow rate of 0.8 mL min⁻¹. The injected volume was 15.0 μ L. Individual PAH identification was carried out by comparison of their retention times with those of their respective standards. External calibrations with PAHs mixed standards, using at least 6 calibration points, were performed. Analytical blanks and standards were analyzed daily and regularly. Each analysis was performed at least in triplicate.

3 RESULTS AND DISCUSSION

3.1 $PM_{2.5}$ concentrations

$PM_{2.5}$ exposure levels of firefighters in the characterized four Portuguese fire stations ranged between 0.05 to 1.04 μ g/m³.

Studies concerning firefighters' exposure to $PM_{2.5}$ are limited, particularly at fire stations. A study conducted in American wildland firefighters at non prescribed forest burn days assessed personal $PM_{2.5}$ concentrations ranging from 9 and 43 μ g/m³ during a work shift of 7 to 12 hours (Adetona et al. 2011). More recently, a study conducted also in American firefighters reported $PM_{2.5}$ concentrations ranging between 18 and 155 μ g/m³ at two firehouses (Baxter et al. 2014). In this work, the determined $PM_{2.5}$ concentrations were well below the values reported by those studies.

Regarding occupational exposure, the National Institute for Occupational Safety and Health (NIOSH) defined a permissible exposure limit of 15 and 5 μ g/m³ for total and respirable particles not otherwise regulated, respectively (NIOSH 2007). Firefighters' personal exposure to $PM_{2.5}$ inside Portuguese fire stations was below those guidelines.

3.2 PAH concentrations

In this study, $PM_{2.5}$ -bound PAHs were evaluated in Portuguese firefighters' personal air. Individual PAHs namely naphthalene, acenaphthene, fluorene, and phenanthrene were detected in all the samples while acenaphthylene, benzo[b+j]fluoranthene, and benzo[g,h,i]perylene were detected in more than 80% of $PM_{2.5}$ filters. Chrysene, dibenz[a,h]anthracene, and indeno[1,2,3-cd]pyrene were detected in less than 15% of the samples. Among the 18 analyzed PAHs, dibenzo[a,l]pyrene was the only compound that was never detected in the firefighters personal air samples.

Overall, total PAH concentrations ranged between 35.8 to 294 ng/m³. Acenaphthylene,

acenaphthene, naphthalene were the most abundant compounds, accounting for almost 88% of the total PAHs. Other predominant PAHs were, by descending order: dibenz[a,h]anthracene, phenanthrene, and benzo[g,h,i]perylene. Together these six PAHs represented 97% of total PAHs.

As described in previous studies (Oliveira et al. 2015; Slezakova et al. 2014), in air PAHs are distributed between gaseous (predominantly compounds with 2–3 rings) and particulate phases (mainly PAHs with 4 or more aromatic rings). At the four Portuguese fire stations, PM_{2.5} indoor fraction presented significantly higher concentrations of acenaphthylene and acenaphthene. No specific occurrences that could justify the high prevalence of these two PAHs in the indoor PM_{2.5} phase were found. A possible explanation to this evidence may be the contribution from some particular emission source inside fire stations.

Several countries have set occupational exposure limits for coal-tar pitch volatiles or particulate PAHs (IARC 2002). Those reference values are typically based on the benzene—or cyclohexane-soluble matter of the particulates present in air which includes benz[a]anthracene, benzo[b]fluoranthene, chrysene, anthracene, benzo[a]pyrene, phenanthrene, acridine or pyrene. The Agency for Toxic Substances and Disease Registry recommended an airborne exposure limit of 0.1 mg/m³ for coal tar pitch volatiles during a working day (ATSDR 2015). Portuguese firefighters' personal exposure to PAHs was well below those limits. Additionally, the American Conference of Governmental Industrial Hygienists states that, for any substance with no numerical threshold limit value (but designated as a suspected human carcinogen), worker exposure by all routes should be carefully controlled to levels as low as possible (ACGIH 2010).

Only two studies were found in the literature regarding firefighters exposure to PAHs at fire stations. Shen et al. (2015) focused on dust samples collected from different vacuum cleaner bags, which were used to routinely clean twenty Californian fire stations. Authors reported total PAHs concentration of 4124 ng/g in dust, being pyrene, benzo[ghi]perylene, fluoranthene, and chrysene the most predominant PAHs (Shen et al. 2015). Baxter et al. (2014) conducted a study in two different metropolitan fire stations located in Cincinnati, USA. Air monitoring was performed in the fire truck bay, equipment storage area, kitchen/common area, and in sleeping quarters during a period of 8-hours on a single day. Among the seventeen screened PAHs, only naphthalene was found in one fire station, with mean concentrations of 9.22 µg/m³ and 9.24 µg/m³ at the kitchen and truck bay, respectively (Baxter et al. 2014). Mean concentration of naphthalene was 9.69 ng/m³ (2.03–14.3 ng/m³) at four Portuguese fire stations which was similar to

those reported by Baxter et al. (2014). The concentrations of naphthalene in Portuguese fire stations were well below the short-term exposure limit of 11 mg/m³ over a period of 15-minutes of exposure designated by the US National Institute for Occupational Safety and Health (NIOSH 2007).

Due to the presence of aromatic rings, PAHs present high toxicity, mutagenicity and carcinogenic characteristics (IARC 2002, 2010). Benzo[a]pyrene, the biomarker of carcinogenicity, was detected in 67% of the samples with concentrations ranging between 6.74×10^{-2} and 1.00 ng/m³ (mean of 0.23 ng/m³) in the indoor air of fire stations. Overall, total concentrations of carcinogenic PAHs ranged from 2.89 to 23.7 ng/m³ (mean of 20.9 ng/m³) and accounted with 12% for the total PAHs.

4 CONCLUSIONS

This work provides relevant information regarding occupational exposure of Portuguese firefighters to PM_{2.5} and PM_{2.5}-bound PAHs in the indoor air of four fire stations. Globally, PM_{2.5} concentrations were well below the Portuguese guideline and as a consequence might not represent significant risks for firefighters health. Total PM_{2.5}-bound PAHs ranged between 35.8 and 294 ng/m³ and total concentrations of carcinogenic PAHs accounted with 12% for the total PAHs. Since some PAHs have been reported to possess reproductive, developmental, hemato-, cardio-, neuro-, and immuno-toxicities in human and laboratory animals (ATSDR 1995), more studies regarding firefighters occupational exposure to PAHs are needed in order to better characterize the respective exposures.

ACKNOWLEDGEMENTS

This work was supported by European Union (FEDER funds through COMPETE) and National Funds (Fundação para a Ciência e Tecnologia) through projects UID/QUI/50006/2013 and UID/EQU/00511/2013-LEPABE, by the FCT/MEC with national funds and co-funded by FEDER in the scope of the P2020 Partnership Agreement. Additional financial support was provided by Fundação para a Ciência e Tecnologia through fellowships SFRH/BD/80113/2011 (M. Oliveira), SFRH/BPD/65722/2009 (K. Slezakova).

REFERENCES

ACGIH 2010. *Documentation for a Recommended BEI of Polycyclic Aromatic Hydrocarbons*. American

- Conference of Governmental Industrial Hygienists, Cincinnati, Ohio, USA.
- Adetona, O., Dunn, K., Hall, D.B., Achtemeier, G., Stock, A. & Naeher, L.P. 2011. Personal PM_{2.5} exposure among wildland firefighters working at prescribed forest burns in Southeastern United States. *Journal of Occupational and Environmental Hygiene* 8:503–511.
- ATSDR 2015. *Environmental Health and Medicine Education—Polycyclic Aromatic Hydrocarbons*. Agency for Toxic Substances and Disease Registry. Agency for Toxic Substances and Disease Registry. Available from: <http://www.atsdr.cdc.gov/csem/csem.asp?csem=13&po=8> (last accessed 16-09-2015).
- ATSDR 1995. *Toxicological Profile for Polycyclic Aromatic Hydrocarbons*, Atlanta: Agency for Toxic Substances and Disease Registry. Available from: <http://www.atsdr.cdc.gov/toxprofiles/tp69.html> (last accessed 16.09.2015).
- Baxter, C.T., Hoffman, J.D., Knipp, M.J., Reponen, T. & Haynes, E.N. 2014. Exposure of firefighters to particulates and polycyclic aromatic hydrocarbons. *Journal of Occupational and Environmental Hygiene* 11:D85–D91.
- Castro, D., Slezakova, K., Oliva-Teles, M.T., Delerue-Matos, C., Alvim-Ferraz, M.C., Morais, S. & Pereira, M.C. 2009. Analysis of polycyclic aromatic hydrocarbons in atmospheric particulate samples by microwave-assisted extraction and liquid chromatography. *Journal of Separation Science* 32:501–510.
- Castro, D., Slezakova, K., Delerue-Matos, C., Alvim-Ferraz, M., Morais, S. & Pereira, M.C. 2011. Polycyclic aromatic hydrocarbons in gas and particulate phases of indoor environment influenced by tobacco smoke: levels, phase distribution and health risks. *Atmospheric Environment* 45: 1799–1808.
- Directive 2004/107/EC, 2005. *Directive of the European Parliament and of the Council relating to arsenic, cadmium, mercury, nickel and polycyclic aromatic hydrocarbons in ambient air*. Official Journal of European Union, L23, 3–16.
- Estrellan, C.R. & Lino, F. 2010. Toxic emissions from open burning. *Chemosphere* 80, 193–207.
- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. 2002. *Some traditional herbal medicines, some mycotoxins, naphthalene and styrene*. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans 82: 1–556.
- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. 2010. *Some non-heterocyclic polycyclic aromatic hydrocarbons and some related exposures*. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans 92:1–853.
- Lemieux, P.M., Lutes, C.C. & Santoianni, D.A. 2004. Emissions of organic air toxics from open burning: a comprehensive review. *Progress in Energy and Combustion Science* 30:1–32.
- Lewtas, J. 2007. Air pollution combustion emissions: characterization of causative agents and mechanisms associated with cancer, reproductive, and cardiovascular effects. *Mutation Research/Reviews in Mutation Research* 637:95–133.
- NIOSH. 2007. *NIOSH Pocket Guide to Chemical Hazards*. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention. National Institute for Occupational Safety and Health. Cincinnati, Ohio.
- Okona-Mensah, K.W., Battershill, J., Boobis, A. & Fielder, R. 2005. An approach to investigating the importance of high potency polycyclic aromatic hydrocarbons (PAHs) in the induction of lung cancer by air pollution. *Food and Chemical Toxicology* 43:1103–1116.
- Oliveira, M., Slezakova, K., Delerue-Matos, C., Pereira, M.C. & Morais, S. 2015. Polycyclic aromatic hydrocarbons: levels and phase distributions in pre-school microenvironment. *Indoor Air* 25(5):557–568.
- Reisen, F., Brown, S. & Cheng, M. 2006. Air toxics in bushfire smoke: firefighter's exposure during prescribed burns. *Forest Ecology and Management* 234:S144.
- Shen, B., Whitehead, T.P., McNeel, S., Brown, F.R., Dhaliwal, J. & Das, R. et al. 2015. High levels of polybrominated diphenyl ethers in vacuum cleaner dust from California Fire Stations. *Environmental Science & Technology* 49:4988–4994.
- Slezakova, K., Castro, D., Delerue-Matos, C., Morais, S. & Pereira, M.C. 2014. Levels and risks of particulate-bound PAHs in indoor air influenced by tobacco smoke: A field measurement. *Environmental Science and Pollution Research* 21:4492–4501.
- USEPA. 2015. *Risk-Based Concentration Table*. United States Environmental Protection Agency. http://www.epa.gov/reg3hwmd/risk/human/rb-concentration_table/usersguide.htm, accessed in October 2015.
- WHO. 2006. *Air Quality Guidelines Global Update 2005*, World Health Organization, Regional Office for Europe, Copenhagen, Denmark, ISBN 92–890–2192–6.
- WHO. 2013. *State of the science of endocrine disrupting chemicals 2012*. United Nations Environment Programme and the World Health Organization, Geneva.