

The effect of a multicomponent training program in elderly's body composition.

António M Monteiro^{1,2}, Emília Alves³, Pedro Forte^{2,3}

¹Instituto Politécnico de Bragança, Bragança, Portugal; ²Research Center in Sports Sciences, Health and Human Development, Portugal; ³Higher Institute of Educational Sciences of the Douro, Penafiel, Portugal;

Introduction

Physical exercise is recommended to maintain the muscular mass in elderlies (Singh, 2002). Among the different types of exercise programs, there is the multicomponent training (Carvalho et al., 2010). The multicomponent training program allows the subjects to improve their physical condition (Strength, resistance, flexibility and balance) (Carvalho et al., 2009). However, there is no consensus about the influence of multicomponent training in body composition. Thus, the aim of this study was to access the multicomponent training effect in elderly's body composition..

Methods

The sample of this study was composed with forty-nine elderlies with 64.39 (\pm 6.33) years old. Among them, 11 were males with 67.45 (\pm 4.93) and 38 were females with 63.50 (\pm 7.47) years old. All procedures carried out in this research were in accordance to the Declaration of Helsinki. A multicomponent training program was applied with the Carvalho et al, (2009). The program had 8 months and the training frequency was three times per week.

The elderly's body composition were assessed with bio-impedance (Tanita, BC-545). The T-test allowed to assess the differences between the pre and post training program in the body composition. The tests were performed with a significant level of 5%.

Results

Table 1 presents the mean and standard deviation (\pm SD) between the two evaluation moments. The statistical significance is also presented in table 1.

Table 1. Mean and standard deviation (\pm SD) of the body composition variables between the pre and post-test.

Variables	Pre-Test	Post-Test	p
	(\pm SD)	(\pm SD)	
Body mass	68.92 (\pm 8.58)	68.51 (\pm 8.69)	*
Fat mass	32.06 (\pm 7.723)	31.55 (\pm 7.840)	0.011
Muscular mass	44.92 (\pm 7.23)	45.14 (\pm 9.28)	0.873
Body water (%)	49.49 (\pm 6.27)	49.67 (\pm 6.37)	0.004
Metabolic rate	1390.02 (\pm 202.67)	1415.02 (\pm 202.84)	0.006
Bone mineral mass	2.24 (\pm 0.48)	2.24 (\pm 0,48)	*
Visceral fat	9.61 (\pm 2.72)	9.35 (\pm 2.61)	**

*p<0.05; **p<0.001

Conclusion

A multicomponent training program in elderlies seem to influence the body composition. The total body mass decreased between the two moments. The bone mineral mass and body water percentage had a significant increased between the two moments. Even more, all the variables improved between the two moments.

References

- Singh, M. A. (2002). Exercise comes of age: rationale and recommendations for a geriatric exercise prescription. *J Gerontol A Biol Sci Med Sci*, 57(5), M262-282.
- Carvalho, J., Marques, E., Soares, J. M., & Mota, J. (2010). Isokinetic strength benefits after 24 weeks of multicomponent exercise training and a combined exercise training in older adults. *Aging Clin Exp Res*.
- Carvalho, M. J., Marques, E., & Mota, J. (2009). Training and detraining effects on functional fitness after a multicomponent training in older women. *Gerontology*, 55(1), 41-48.