


BIOTEC'98 BOOK OF ABSTRACTS



IV Iberian Congress on Biotechnology
I Ibero-American Meeting on Biotechnology

BIOTEC'98

BOOK OF ABSTRACTS

Edited by

MANUEL MOTA and EUGÉNIO C. FERREIRA

Biodegradation of Textile Azo Dyes by *P. chrysosporium*

Isabel C. Ferreira, Isabel M. Santos, M^a João Queiroz* and Nelson Lima
Instituto de Biotecnologia e Química Fina
Universidade do Minho
Campus de Gualtar
4710 Braga-Portugal

Key-words: Azo dyes, *P. chrysosporium*, biodegradation, decolorization, effluent treatment.

Azo dyes are used extensively in the textile and dyestuff industries and effluents from these industrial processes are usually resistant to biological treatment.

Textile *azo* dyes with bioaccessible groups such as guaiacol and 2,6-dimethoxyphenol, for lignin-degrading fungus as *P. chrysosporium* were synthesised (Fig. 1).




Fig. 1

Some of these dyes were studied in order to compare the biodegradation ability related to their chemical structure, when cultured with *P. chrysosporium*.

The degradation of dyes was followed by UV-Visible spectrophotometric assays studying the degree of decolorization by the decrease of the absorbance in the maximum wavelength of the dye (Fig. 2) or by the possible formation of new products and by biomass production.




Fig. 2

Thanks are due to IBQF/UM (Praxis XXI) and to Praxis XXI 2/2.1/QUI/44/94 for financial support and I.M. Santos support by a grant Praxis XXI/BD/9120/96.