




*Food,
Timber,
Biomass &
Energy in
Europe*

**Cuneo, Italy
13-16 October**

Castanea 2009

**1st European Congress on Chestnut
5° Convegno Nazionale Castagno**

ABSTRACTS



COMITATO SCIENTIFICO / SCIENTIFIC COMMITTEE

Giancarlo Bounous	Presidente / Convener Università di Torino
Alberto Alma	Università di Torino
Emilio Amorini	CRA Istituto Sperimentale per la Selvicoltura, Arezzo
Sanzio Baldini	Università della Tuscia, Viterbo
Roberto Botta	Università di Torino
Tiziano Caruso	Università di Palermo
Orazio Ciancio	Università di Firenze
Alessandro Crosetti	Università di Torino
Carlo Fideghelli	CRA Istituto Sperimentale per la Frutticoltura, Roma
Raffaello Giannini	Università di Firenze
Vittorino Novello	Università di Torino
Giovanni Nicolotti	Università di Torino
Cristiana Peano	Università di Torino
Davide Pettenella	Università di Padova
Pietro Piccarolo	Università di Torino
Mario Pividori	Università di Padova
Carlo Pirazzoli	Università di Bologna
Giuseppe Scarascia Mugnozza	Università della Tuscia, Viterbo
Giacomo Tamietti	Università di Torino
Tullio Turchetti	Consiglio Nazionale delle Ricerche, Firenze
Andrea Vannini	Università della Tuscia, Viterbo
Milan Bolvansky	Slovak Republic
Mihai Botu	Romania
Svetla	Bulgaria
Bratanova-Doncheva	
Marco Conedera	Switzerland
Stephanos Diamandis	Greece
Victor Galan Sauco	Spain
Juan F. Gallardo Lancho	Spain
Bernard Hennion	France
Afonso Martins	Portugal
Ibrahim Mujic	Croatia
Lorenzo Santiago Pereira	Spain
Mikhail Pridnya	Russia
Laszlo Radocz	Hungary
Arif Soyly	Turkey
Anita Solar	Slovenia

SEGRETERIA SCIENTIFICA / SCIENTIFIC SECRETARY

Gabriele Loris Beccaro	<i>Dipartimento di Colture Arboree</i>
Maria Gabriella Mellano	<i>Università di Torino</i>
Sara Canterino	



Castanea 2009

Altino Choupina ^(1,2), Sofia Meirinho ⁽¹⁾, Marisa Carvalho ⁽¹⁾, Lurdes Jorge ⁽¹⁾, M. João Sousa ⁽¹⁾ and Alfredo Cravador ⁽³⁾

⁽¹⁾ Instituto Politécnico de Bragança, Escola Superior Agrária, Campus de Santa Apolónia - Apartado 1172, 5301-854 BRAGANÇA, PORTUGAL

⁽²⁾ CIMO Campus de Santa. Apolónia - Apartado 1172, 5301-854 BRAGANÇA, PORTUGAL

⁽³⁾ Universidade do Algarve (UAIG), Campus da Penha, Estrada da Penha. 8005-139 FARO, PORTUGAL

INTRODUCTION

Phytophthora species are the causal agents of many serious plant diseases. They secrete large amounts of elicitors, a group of unique highly conserved proteins that are able to induce hypersensitive response (HR) and enhance plant defense responses in a systemic acquired resistance (SAR) manner against infection by different pathogens. *Phytophthora cinnamomi*, one of the most destructive species of *Phytophthora* genus is the causal agent of *Castanea sativa* ink disease, and has been associated with the decline of several forest, ornamental and fruit trees and shrubs, causing enormous economic losses worldwide.

We briefly describe some of the proteins involved in *P. cinnamomi* infection mechanisms: a transglutaminase (TGase, induction of defense responses and disease-like symptoms), a glucanase inhibitor protein (GIP, causing suppression of host defense responses) and a necrosis-inducing protein 1 (NPP1).

MATERIALS AND METHODS

All genes referred in this study were obtained from *P. cinnamomi* strain Pr120 genomic DNA. Elucidation of complete gene nucleotide sequences of TGase, GIP and NPP1 were achieved by High-Efficiency Thermal Asymmetric Interlaced PCR (HE-TAIL PCR), a method described by Michiels *et al.* (2003). DNA sequencing was performed using an ABI 373 automated sequencer. The open reading frames (ORF) of *P. cinnamomi* genes were identified by BioEdit program and submitted to EMBL databases (GIP-accession number CAJ90742.1; *P. cinnamomi* transglu gene for transglutaminase elicitor precursor, accession AM403129; *P. cinnamomi* npp1 gene for necrosis-inducing protein - accession AM403130). Nucleotide and amino acid sequences were analyzed using FASTA programs from EMBL databases. ClustalW2 (Larkin *et al.*, 2007) was used to align the *Phytophthora* genus sequences.

RESULTS

The translated ORF of *P. cinnamomi* GIP codifies a 269aa protein, with a predict Mw of 28,8KDa. Scanning against protein search databases revealed that *P. cinnamomi* GIP are a serine protease, with a trypsin domain profile. A characteristic feature of Ser proteases is to have a catalytic triad charge relay system, with residues of H, D and S in that order along the sequence, essential for the proteolytic function. In Figure 1 are shown the multiple alignment of various sequences who showed great homology with *P. cinnamomi* GIP, including another GIPs of *Phytophthora* genus, and a serine protease and a trypsin protease from *P. infestans*. GIPs have in common the fact that none of them have an intact catalytic triad, like other serine proteases, although they share with them several stretches of amino acids and motifs that are highly conserved. Thus, in all *Phytophthora* GIPs, there are substitutions in residues of the catalytic triad: H-79→A,S,T,I,M-79 (in *P. cinnamomi*: S-79), D-128→N-128 (only in *P. cinnamomi* and *P. sojae* GIP2, and Ser-217→T-217, in all *Phytophthora* GIPs. Therefore, GIPs are proteolytically inactive, referred as serine protease homologs, and presumably function as host-enzyme inhibitors. It can be hypothesized that a major role for GIP is to suppress the release of glucan elicitors during *Phytophthora sp.* infections, thereby reducing the effectiveness of the plant host's surveillance system (Rose *et al.*, 2002).




FIGURE 1 – Multiple sequence alignment of GIP and GIP-like genes from *Phytophthora sp.*

B0B0H5_PHYCIN – putative GIP from *P. cinnamomi*; Q945U0_9STRA – GIP1 from *P. sojae*; B1AC90_PHYIN – GIP1 from *P. infestans*; Q945T9_9STRA – GIP2 from *P. sojae*; B1AC87_PHYIN – GIP2 from *P. infestans*; Q945T8_9STRA – GIP2 from *P. sojae*; B1AC88_PHYIN – GIP3 from *P. infestans*; Q2M411_PHYIN – trypsin protease GIP-like; B1AC89_PHYIN – GIP4 from *P. infestans*; B1AC86_PHYIN – serine protease from *P. infestans*; Q2M412_PHYIN – trypsin protease GIP-like.

P. cinnamomi transglutaminase is a protein with 533 aa with a predict Mw of 57,7KDa. *Phytophthora sp.* TGases are even more closely related amongst them than GIPs, as shown in Figure 2.

P. cinnamomi necrosis-inducing protein1 has 256 aa with a predict Mw of 29,0KDa. Scanning against protein search databases (data not shown) revealed that sequences who showed greater homology with *P. cinnamomi* NPP1 were two NPP1 proteins of *P. infestans* (Q2M430_PHYIN; Q2M429_PHYIN). The phylogram showing the more closely related sequences are shown in Figure 3.




FIGURE 2 – Multiple sequence alignment of TGase genes from *Phytophthora sp.* B0B0Q5_PHYCI - *P. cinnamomi* transglutaminase elicitor; Q6Q475_9STRA – *P. sojae* transglutaminase elicitor; Q01928_PHYME – *P. megasperma* glycoprotein elicitor; Q6XDM3_PHYIN - *P. infestans* transglutaminase elicitor M81C; Q6XDM2_PHYIN - *P. infestans* transglutaminase elicitor M81D)




FIGURE 3 – Phylogram of most closely related *P. cinnamomi* NPP1 protein. B0B0Q6_PHYCI - *P. cinnamomi* NPP1; Q8LKL0_9STRA – *P. sojae* necrosis-inducing-like protein; Q9AT28_PHYPR – *P. parasitica* necrosis-inducing-like protein; Q2M430_PHYIN - *P. infestans* NPP-like protein; Q2M429_PHYIN - *P. infestans* NPP-like protein.

Further studies, including plant-pathogen phenotypic interactions are needed to understand how each individual factor can affect pathogenesis mechanisms of *P. cinnamomi*, and how this knowledge can be used for control ink disease and other *Phytophthora sp.* diseases.

REFERENCES

- Larkin, M., Blackshields, G., Brown, N., Chenna, R., McGettigan, P., McWilliam, H., Valentin, F., Wallace, I., Wilm, A., Lopez, R., Thompson, J., Gibson, T. & Higgins D., 2007. Bioinformatics, 23(21): 2947-2948.
Michiels, A., Tucker, M., Van Den Ende, W. & Van Laere, A., 2003. Chromosomal Walking of Flanking Regions From Short Known Sequences in GC-Rich Plant Genomic DNA. *Plant Molecular Biology Reporter*, 21:295-302.
Rose, J., Ham, K., Darvill, A. & Albersheim, P., 2002. Molecular cloning and characterization of glucanase inhibitor proteins: coevolution of a counterdefense mechanism by plant pathogens. *Plant Cell*, 14:1329-1345.

ACKNOWLEDGEMENTS: The Project COMBATINTA/SP2.P11/02 Interreg IIIA – Cross-Border Cooperation Spain-Portugal, financed by The European Regional Development Fund, and the Project “Identification, characterization and role of molecular factors associated with the mechanisms of infection of Fagaceae species by *Phytophthora cinnamomi*” (PTDC/AGR-AAM/67628/2006) FCT, supported this work.