

IV INTERNATIONAL CONFERENCE
ON ENVIRONMENTAL, INDUSTRIAL
AND APPLIED MICROBIOLOGY

BioMicroWorld 2011

• Torremolinos, Málaga, Spain / 14-16 September 2011

<http://www.formatex.org/biomicroworld2011>

Book of Abstracts



IV International Conference on Environmental, Industrial and Applied Microbiology

BioMicroWorld2011

Torremolinos (Spain), 14-16 September 2011

<http://www.formatex.org/biomicroworld2011>

BioMicroWorld2011 Conference is a multi-disciplinary forum that brings together active researches involved in environmental, industrial and applied microbiology with the aim of communicating current research priorities and progress in the field, and identifying new approaches and research opportunities in applied microbiology.

BioMicroWorld consolidates a series of conferences whose current edition has received about 950 abstracts to be considered for presentation and has attracted nearly 575 researchers from more than 60 countries.

TOPICS

- Agriculture, Soil, Forest Microbiology
- Environmental, Marine, Aquatic Microbiology, Geomicrobiology
- Food Microbiology
- Industrial Microbiology - Future Bioindustries
- Medical Microbiology - Pharmaceutical Microbiology - Antimicrobial agents and chemotherapy
- Methods - Quantitative Models and Bioinformatics in Microbiology - Technology development
- Microbial Physiology, Metabolism and Gene Expression
- Biodegradation and Bioremediation
- Biotechnologically Relevant Enzymes and Proteins
- Biofilms

ORGANIZING COMMITTEE

- A. Méndez-Vilas**, Formatex Research Center, Badajoz, Spain (General Coordinator)
J. A. Mesa González, Formatex Research Center, Badajoz, Spain (Secretariat)
A. Solano Martín, Formatex Research Center, Badajoz, Spain (Secretariat)
J. Mesa González, Formatex Research Center, Badajoz, Spain
A. Agudo Rodríguez, Formatex Research Center, Badajoz, Spain

SCIENTIFIC ADVISORY COMMITTEE

- Ece Karatan**, Appalachian State University, USA
Douglas Weibel, University of Wisconsin-Madison, USA
Anil Kumar Puniya, National Dairy Research Institute, India
Jan Michiels, Centre of Microbial and Plant Genetics, K.U.Leuven, Belgium
Rosario Muñoz, Institute of Industrial Fermentation, Spain
Vladimir Jiranek, The University of Adelaide, Australia
Ibrahim Banat, University of Ulster, United Kingdom
Hongkai Wu, Hong Kong University of Science and Technology, Hong Kong
Jan Nesvera, Institute of Microbiology, Academy of Sciences of the Czech Republic, Czech Republic
Filip Boyen, Ghent University, Belgium
Ramesh C Kuhad, University of Delhi South Campus, India
Essaid A. Barka, University of Reims, France
Jose Luis Martinez, National Center for Biotechnology, Spain
Joonhong Park, Yonsei University, Republic of Korea
Marcia Nitschke, University of Sao Paulo, Brazil
Raeid M. M. Abed, Sultan Qaboos University, Sultanate of Oman
James Gomes, Indian Institute of Technology Delhi, India
Petr Baldrian, Institute of Microbiology ASCR, Czech Republic
Alban Ramette, Max Planck Institute for Marine Microbiology, Germany
Matti Karp, Tampere University of Technology, Finland
Joseph Kreit, Mohammed V University, Morocco
Igor B. Zhulin, University of Tennessee - Oak Ridge National Laboratory, USA
Yves Blache, University of the South, Toulon-Var, France
Manuel Simões, University of Porto, Porto, Portugal
Badal Saha, National Center for Agricultural Utilization Research USDA-ARS, USA
David A McDowell, University of Ulster, United Kingdom
Carsten S. Jacobsen, Geological Survey of Denmark and Greenland, GEUS, Denmark
Hans van Veen, Netherlands Institute of Ecology, Heteren, The Netherlands
Kaarina Sivonen, Helsinki University, Finland
Bruce Maguire, Pfizer Global Research and Development, USA
Juan José Valdez Alarcón, Michoacana University of Saint Nicolas Hidalgo, Mexico
Rodney M. Donlan, Centers for Disease Control and Prevention, USA
Jean-Luc Jany, Laboratory of Biodiversity and Microbial Ecology, University of Brest, France
Pilar García, Asturias Dairy Products Institute, Spain
Pei-Yuan Qian, Hong Kong University of Science and Technology, Hong Kong
Carme Plumed-Ferrer, University of Eastern Finland, Finland
Eduardo Diaz, Centre for Biological Research CIB (CSIC), Spain
Ivan Berg, University of Freiburg, Germany
Marc Solioz, University of Berne, Switzerland
Brian B. McSpadden Gardener, The Ohio State University, USA
Tino Krell, Experimental Station of El Zaidin (CSIC), Spain
Thomas Maskow, Helmholtz Centre for Environmental Research - UFZ, Germany
Vasu Appanna, Laurentian University, Canada
Manfredo J. Seufferheld, University of Illinois at Urbana-Champaign, USA

Effect of nitrogen supplementation on yeast fermentation performance and mead quality

A. P. Pereira^{1,2}, A. Mendes-Ferreira², J. M. Oliveira³, L. M. Estevinho¹ and A. Mendes-Faia²

¹CIMO, Centro de Investigação de Montanha, Escola Superior Agrária, Instituto Politécnico de Bragança, Campus de Santa Apolónia - Apartado 1172, 5301-855 Bragança, Portugal

²IBB, Institute for Biotechnology and Bioengineering, Centre of Genomic and Biotechnology, Universidade de Trás-os-Montes e Alto Douro, Apartado 1013, 5001-801 Vila Real, Portugal

³IBB, Institute for Biotechnology and Bioengineering, Centre of Biological Engineering, Universidade do Minho, 4710-057 Braga, Portugal.

Mead is a traditional drink, containing 8-18% (v/v) of ethanol, which results from the alcoholic fermentation of diluted honey performed by yeasts. However, when it is produced in a homemade way, mead producers find several problems, namely, the lack of uniformity in the final product, slow or premature fermentations arrest, and the production of "off-flavours" by the yeasts. These problems could be due to several factors, including lack of essential nutrients such as a deficiency in available nitrogen. Additionally, it has been reported that mead fermentation is a time-consuming process, often taking several months to complete, depending on the type of honey, yeast strain and honey-must composition. Since mead production is a time-consuming process, to make its production viable it is necessary to reduce the fermentation time while producing an end product of quality. Thus, the aim of this study was to evaluate the effect of nitrogen addition to honey-must on two active dry wine yeasts (ADWY) fermentation performance, as well as on the mead composition and volatile aroma compounds production.

Honey must was prepared according a recipe developed in our laboratory, and supplemented with potassium tartrate and pH adjusted to 3.7 with malic acid. Then to study the effect of nitrogen addition a part of honey-must was adjusted with diammonium phosphate (DAP) to achieve the concentration of nitrogen required by yeast to complete alcoholic fermentation. The honey-musts were inoculated in order to obtain a pitching rate of 1×10^7 viable cells/ml. Several parameters were determined during the fermentation to evaluate the effect of nitrogen addition on yeast growth, fermentation profile, mead composition and mead aromatic profile. For this study as biological material were selected the ADWY *Saccharomyces cerevisiae* Lalvin QA23 and Lalvin ICV D47.

The supplementation of honey-must with DAP reduced fermentation length in approximately seven days, however sugars were not fully consumed, suggesting that other factors could be interfering with yeast growth. Furthermore, it was verified that for both yeasts the specific growth rate and final biomass were higher in musts supplemented with DAP. Mead final composition was similar in the two experimental conditions, however, even in the honey-must to which DAP was not added about 25 mg/L of assimilable nitrogen remained at the end of fermentation. Some fermentative aroma compounds which contribute to the sensorial quality of mead, including alcohols, fatty acid ethyl esters, acetates, volatile phenols and volatile fatty acids, were identified and quantified. Global analysis of volatile profile revealed that the concentration of fatty acid ethyl esters and volatile phenols was higher in meads supplemented with DAP. The concentration of volatile phenols was below their perception threshold but the levels of acetate and ethyl esters could contribute to enhance fruity character in meads produced. These results are very useful for optimise the mead production and improving its quality.

Keywords mead; assimilable nitrogen, yeast performance, aromatic profile

A.P.P. was recipient of a PhD grant from FCT (SFRH/BD/45820/2008).