




FACULTY OF SCIENCE
UNIVERSITY OF MINHO



3rd Symposium on **Medicinal Chemistry** of University of Minho

May 26th, 2017

School of Science, Chemistry Department

University of Minho, Campus of Gualtar

Scientific Committee

Fernanda Proença - Department of Chemistry, University of Minho

Isabel Ferreira - Polytechnical Institute Bragança

M. Alice Carvalho - Department of Chemistry, University of Minho

A. Gil Fortes - Department of Chemistry, University of Minho

Luis S. Monteiro - Department of Chemistry, University of Minho

Organizing Committee

Maria Alice Carvalho

António Gil Fortes

Luis Monteiro

Ana Patrícia Lourenço (MSc student of Medicinal Chemistry)

Ana Rita Abreu (MSc student of Medicinal Chemistry)

Ana Rita Araújo (MSc student of Medicinal Chemistry)

André Carvalho (MSc student of Medicinal Chemistry)

David Freitas (MSc student of Medicinal Chemistry)

Diogo Lopes (MSc student of Medicinal Chemistry)

Inês Machado (MSc student of Medicinal Chemistry)

Joana Filipa Parente (MSc student of Medicinal Chemistry)

Juliana Rocha (MSc student of Medicinal Chemistry)

Mónica Fernandes (MSc student of Medicinal Chemistry)

Pedro Figueiredo (MSc student of Medicinal Chemistry)

Posters

- P01 Phenolic Profile, Tocopherols and Antioxidant Activity of Watercress** (41)
José Pinela
- P02 Serine HydroxymethylTransferase: Inhibit it or Improve it?** (42)
H. S. Fernandes
- P03 Organic acids, tocopherols and antioxidant activity of white and black garlic samples** (43)
Joana Botas
- P04 From the raw material to your headache** (44)
Iva Costa
- P05 Are you confident about your Chromatographic Result?** (45)
Katalin Kováts
- P06 Improving the bioactivity of medicinal plants using gamma radiation: Thymus vulgaris L. infusions as a case-study.** (46)
Eliana Pereira
- P07 The role of chamomile phenolic compounds in the development of dairy functional foods** (47)
C. Caleja
- P08 Rosa canina L. as new food ingredient an source of bioactive compounds** (48)
T.C. S. P. Pires
- P09 Synthesis of *N*-catechoyl phenylalanine and dehydrophenylalanine derivatives** (49)
I. Machado
- P10 Synthesis of the methyl esters of *N*-phenoyl and *N*-catechoyl dehydroalanine and dehydrophenylalanine** (50)
S. Oliveira
- P11 Synthesis of phenolic and catecholic amino acids by the modified multicomponent Ugi reaction** (51)
L.S. Monteiro
- P12 Synthesis of phenolic and catecholic amino acids by Michael addition to a dehydroalanine derivative** (52)
J. Parente
- P13 Theoretical and computational studies addressed to the stereo and regio- selective synthesis of polyhydroxylated sugar-like compounds** (53)
J. F. Rocha
- P14 *Pseudofistulina radicata* (Schwein) Burds: the first study focused on its antioxidant activity and bioactive compounds** (54)
Blanca N. Castaneda
- P15 Synthesis of a low molecular weight peptide hydrogelator based on dehydroamino acids** (55)
Rita L. Araújo
- P16 An efficient method for dehydration of 6-carbamoylpurines** (56)
Mónica C. S. Fernandes

Rosa canina L. as new food ingredient an source of bioactive compounds

T.C. S. P. Pires^a, M.I. Dias^a, L. Barros^a, R.C. Calheta^a, M.B. P.P. Oliveira^b, C. Santos-Buelga^c,
I.C.F.R. Ferreira^a

^a Mountain Research Center (CIMO), ESA, Polytechnic Institute of Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal; iferreira@ipb.pt

^bREQUIMTE/LAQV, Science Chemical Department, Faculty of Pharmacy of University of Porto, Rua Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal.

^cGrupo de Investigación en Polifenoles (GIP-USAL), Facultad de Farmacia, Universidad de Salamanca, Campus Miguel de Unamuno s/n, 37007 Salamanca, España

Sustainable food options are increasing among the most concerned consumers, in which they seek to combine new ingredients with potential health benefits [1]. Edible flowers provide new colors, textures and vibrancy to any dish, and apart from the "glam" factor, they can provide bioactive compounds [2]. In the present work, the edible petals of *Rosa canina* L., gently supplied by the RBR Foods Company (Portugal), were tested for their cytotoxic activity against tumor cell lines. The assays were conducted in the hydromethanolic extract and in the lyophilized infusion, being both of them re-dissolved in water in order to obtain stock solutions at 100 mg/mL for successive dilutions until determination of GI₅₀ values (concentration that inhibits 50% of the net cell growth). Four human tumor cell lines were tested: MCF-7 (breast adenocarcinoma), NCIH460 (non-small cell lung cancer), HeLa (cervical carcinoma) and HepG2 (hepatocellular carcinoma). A non-tumour primary cells' culture (assigned as PLP2) was also prepared from a freshly harvested porcine liver, and used in the bioassays. Ellipticine was used as the positive control. While the hydromethanolic extract inhibited the growth of HeLa (308 µg/mL), the lyophilized infusion inhibited MCF-7 (377 µg/mL), respectively. Both extracts presented cytotoxicity for HepG2 (297 µg/mL and 315 µg/mL for hydromethanolic extract and infusion, respectively). None of the extracts showed toxicity towards PLP2. After chemical characterization of the extracts, flavonoids were the only phenolic compounds present, being quercetin-3-O-glucoside and kaempferol-3-O-glucoside the major compounds present in both extracts. These results showed that *R. canina* edible flowers are sources of bioactive compounds related with cytotoxic properties in human tumor cell lines, highlighting their applicability potential in nutraceutical and pharmaceutical fields.

Acknowledgments:

The authors are grateful to the Foundation for Science and Technology (FCT, Portugal) and FEDER under Programme PT2020 for financial support to CIMO (UID/AGR/00690/2013). L. Barros (SFRH/BPD/107855/2015) and M.I. Dias (SFRH/BD/84485/2012) grants.

References:

1. Pires, T. C. S. P. et al., *Food Chem.* **2017**, *220*, 337–343.
2. Lara-Cortés, E. et al., *Rev. Chapingo Ser. Hortic.* **2014**, *101*–116.