


Atención Primaria

www.elsevier.es/ap



PARALLEL SESSIONS: ORAL COMMUNICATIONS

3rd World Congress of Health Research

Viseu - Portugal, 29-30 September 2016

1. ADULT HEALTH

CHARACTERIZATION OF PATIENTS UNDERGOING NONINVASIVE VENTILATION ADMITTED IN UNIT INTERMEDIATE CARE

Matilde Martins^a, Patrícia Ribas^b, Joana Sousa^c, Andreia Cunha^c, Norberto Silva^d, Teresa Correia^a

^aHealth School of Polytechnic Institute of Bragança, Sports Sciences, Health Sciences and Human Development, Portugal; ^bClinic of Arrifana, Portugal; ^cHealth School of Polytechnic Institute of Bragança, Portugal; ^dHealth School of Polytechnic Institute of Bragança, Unit of the Northeast Local Health Bragança, Portugal.

Contact details: matildemartins@ipb.pt

Introduction: Non-invasive ventilation (NIV) is the application of a ventilatory support without resorting to invasive methods. Today it's considered a credible therapeutic option, with enough scientific evidence to support its application in various situations and clinical settings related to the treatment of acute respiratory disease, as well as chronic respiratory disease.

Objectives: Characterize patients undergoing NIV admitted in Unit Intermediate Care (ICU) in the period from October 1st 2015 to June 30th 2016.

Methods: Prospective study conducted in ICU between October 2015 and June 2016. In this study were included all patients hospitalized in this unit (ICU) and in that time period a sample of 57 participants was obtained. As data collection instruments we used a questionnaire for sociodemographic and clinical data and the Braden scale.

Results: Participants were mostly male 38 (66.7%), the average age 69.5 ± 11.3 years, ranging between 43 and 92 years. They weighed on average 76.6 kg (52 and 150), with an average body mass index of 28.5 kg/m² (20 to 58.5). With skin intact 28 (49.1%) with abnormal perfusion 12 (21.1%), with altered sensitivity 11 (19.3%) and a high risk of ulcer on the scale of Braden 37 (65%). The admission diagnosis was respiratory failure 33 (57.3%) and had different backgrounds. We used reused mask 53 (93.0%), the average time of NIV was 7.1 days (1-28), 4.8 days of hospitalization (1-18) and an average of 7.8 IPAP pressure. 11 (19.3%) of the participants developed face ulcer pressure.

Conclusions: The NIV is used in patients with advanced age, obesity, respiratory failure and high risk of face ulcer development.

Keywords: Patients. Noninvasive ventilation.

TELEPHONE SUPPORT LINE OF RHEUMATOLOGY DEPARTMENT: A 4.5 YEARS EXPERIENCE

Eduardo Santos^a, Andréa Marques^b

^aAssociate Research at the Portugal Centre for Evidence-Based Practice (PCEBP): a Collaborating Centre of the Joanna Briggs Institute - Health Sciences Research Unit: Nursing (UICISA: E), Nursing School of Coimbra (ESENFC); ^bRheumatology department, Centro Hospitalar e Universitário de Coimbra, EPE.

Contact details: ejf.santos87@gmail.com

Introduction: In many health services in developed countries, the telephone support has been used as an innovative approach to providing care and answering questions by nurses, developing especially in specific areas such as rheumatology (Brown et al., 2006).

Objectives: To analyze the profile of users and the main reasons of patients that uses the Telephone Support Line of Day Hospital and early arthritis consultation of the CHUC Rheumatology Department, EPE.

Methods: This is a retrospective descriptive-correlational study with a convenience sample of 448 calls. For continuous variables we used the t-student test, for dichotomous variables the χ^2 test and, finally, we performed a cluster analysis by the TwoStep Cluster method through the log-likelihood distance.

Results: Users have an average age of 44.8 years being predominantly male (58%). The cluster analysis allowed to create three groups whose profile shows that the cluster 1 (n = 96/21.4%) are formed by "older" adults that need to clarify mainly doubts about the results of auxiliary examinations tests (AET)/prescriptions and side effects/complications; the cluster 2 (n = 232/51.8%) are formed by young adults that call to change appointments and mainly to clarify doubts on the results of AET/prescriptions; finally, the cluster 3 (n = 120/26.8%) are formed by "middle age" adults that want to clarify doubts about medication and disease management.

Conclusions: Incoming calls cover a wide range of ages and all kinds of rheumatic diseases. This study allows us to identify in which groups it is necessary to make a more detailed educational