



# BOOK OF ABSTRACTS

**TITLE**

Book of Abstracts of the XX EuroFoodChem Congress

**EDITORS**

M. Beatriz P.P. Oliveira, Joana S. Amaral, Manuel A. Coimbra

**EDITION**

Sociedade Portuguesa de Química  
Av. Da República, 45 – 3º Esq  
1050-187 Lisboa – Portugal

**DATE**

June 2019

**ISBN**

ISBN 978-989-8124-26-5



@ Sociedade Portuguesa de Química  
All rights reserved.

The editors state that the content of scientific abstracts is of the responsibility of their respective authors.

## XX EUROFOODCHEM CONGRESS

### Scientific Committee

M. Beatriz Oliveira (Portugal)  
Manuel A. Coimbra (Portugal)  
Marco Arlorio (Italy) – Chair, EuChemS FCD  
Joana Amaral (Portugal) – Secretary, FCD-EuChemS  
Michael Murkovic (Austria) – Treasurer, EuChemS FCD  
Hans-Jacob Skarpeid (Norway)  
Juana Frias (Spain)  
Livia Simon Sarkadi (Hungary)  
Tanja Dcirkovic Velickovic (Serbia)  
Vieno Piironen (Finland)  
Vincenzo Fogliano (The Netherlands)

### Organizing Committee

M. Beatriz Oliveira (Portugal) – University of Porto  
Manuel A. Coimbra (Portugal) – University of Aveiro  
Joana Amaral (Portugal) – Polytechnic Institute of Bragança, FCD-EuChemS

**Congress organized under the auspices of the Food Chemistry Division of the European Chemical Society (FCD-EuChemS) and the Portuguese Chemical Society (SPQ).**



## SPONSORSHIP & SUPPORT

### INSTITUCIONAL



### SPONSORS



Instituto dos Vinhos do Douro e do Porto, I. P.



## Monitoring physicochemical and sensory attributes during debittering of stoned green olives

Ítala M.G. Marx<sup>a,b</sup>, Nuno Rodrigues<sup>a</sup>, Ana C.A. Veloso<sup>c,d</sup>, José A. Pereira<sup>a</sup>, Susana Casal<sup>b</sup>, António M. Peres<sup>a,e,\*</sup>

<sup>a</sup>Centro de Investigação de Montanha, Instituto Politécnico de Bragança, Bragança, Portugal

<sup>b</sup>REQUIMTE, Faculdade de Farmácia da Universidade do Porto, Porto, Portugal


<sup>c</sup>Instituto Politécnico de Coimbra, ISEC, DEQB, Coimbra, Portugal

<sup>d</sup>CEB - Centre of Biological Engineering, University of Minho, Braga, Portugal

<sup>e</sup>Laboratory of Separation and Reaction Engineering - Laboratory of Catalysis and Materials ESA, Instituto Politécnico de Bragança, Bragança, Portugal

peres@ipb.pt

Stoned green olives are traditional table olives produced in the Northeast of Portugal. They are highly appreciated due to their organoleptic characteristics and levels of complex phenols endowed with strong antioxidant activity, having a significant agro-economic relevance. During the natural debittering process, stoned olives are immersed in water, which is changed (each 1-2 days), leading to a reduction of the initial bitterness and increase of sweetness, turning these green olive edible [1]. Monitoring total phenols contents, the bitterness index as well as the basic sensorial attributes through this washing stage is very important. In this work, the debittering process of 110 samples of stoned green olives (*cvs.* Cobrançosa and Negrinha de Freixo), was monitoring during 20 days. After each debittering-washing time-period (water changed each 2-days) bitter, pungent and sweet intensities sensations were assessed by trained panelists. Furthermore, the bitterness index and total phenols contents were also evaluated spectrophotometrically. Figure 1 shows the trends observed for the sensory attributes and chemical parameters along the debittering-washing process. From the boxplots (Figure 1), it is clear that bitterness index and total phenols contents significantly decreased with the time, which could be directly related to the observed decrease of the bitter taste intensity (for both olive cultivars) while the decrease of the pungent sensation could be related with the decrease of the total phenols contents, mainly influenced by the increase of the sweet sensation. Furthermore, as can be inferred, the initial bitterness and pungency intensities of *cv.* Cobrançosa olives were higher than those from *cv.* Negrinha de Freixo. Nevertheless, after 16-20 days of debittering-washing steps, the bitter and pungent intensities perceived by the trained panelists, for both olive cultivars, were of the same order of magnitude, showing that the debittering procedure adopted was successful and technologically consistent.



**Figure 1.** Boxplots showing the time evolution of (A) bitter, (B) pungent and (C) sweet as well as of (D) bitterness index (g Oleuropein/ kg) and (E) total phenols contents (g GAE/ kg) of the stoned green olives (*cvs.* Cobrançosa and Negrinha de Freixo) during the 20 days of debittering-washing process.

**Acknowledgments:** This research was financially supported by Associate Laboratory LSRE-LCM - UID/EQU/50020/2019 - funded by national funds through FCT/MCTES (PIDDAC) and by Project UID/BIO/04469/2013-CEB, and strategic project PEst-OE/AGR/UI0690/2014-CIMO all funded by European Regional Development Fund (ERDF) through COMPETE2020-Programa Operacional Competitividade e Internacionalização (POCI)-and by national funds through FCT-Fundação para a Ciência e a Tecnologia. Ítala M.G. Marx acknowledges Fundação para a Ciência e a Tecnologia (FCT) through FCT PhD reference: (SFRH/BD/137283/2018).

### References:

[1] A. Sousa, I. Ferreira, R. Calhelha, P. Andrade, P. Valentão, R. Seabra, L. Estevinho, A. Bento, J. Pereira. *Bioorganic & Medicinal Chemistry*, 14, (2006) 8533-8538.