

# Micro Biotec'13

PORTUGUESE CONGRESS OF  
MICROBIOLOGY AND BIOTECHNOLOGY

6<sup>th</sup> - 8<sup>th</sup> December | Aveiro Portugal

## Abstracts Book



## Presidentes das Sociedades

SPM Isabel Sá Correia  
SPBT José Teixeira

## Comissão Organizadora

António C.M. Correia (DBUA) Co-chair  
Manuel A. Coimbra (DQUA) Co-chair  
Anabela Pereira (DBUA)  
Artur Alves (DBUA)  
Cláudia Nunes (DQUA)  
Cláudia Passos (DQUA)  
Dulce Helena Teixeira (DQUA)  
Elisabete Coelho (DQUA)  
Maria Ângela Cunha (DBUA)  
Sónia Mendo (DBUA)

## Comissão Científica

António Correia (UA)	João Mano (UM)
Arsénio Fialho (IST, IBB)	João Queiroz (UBI)
Ascensão Reis (UNL)	Joaquim Cabral (IST)
Cecília Arraiano (ITQB/UNL)	Jorge Pedrosa (UM,ICVS)
Cecília Leão (UM, ICVS)	José Teixeira (UM)
Cecília Roque (UNL)	Luísa Peixe (UP, FF)
Célia Manaia (ESB, UCP)	Madalena Alves (UM)
Elsa Anes (FF, UL/IMM)	Manuel Carrondo (FCT/UNL)
Eugénio Ferreira (UM)	Manuel Mota (UM)
Francisco Gírio (LNEG)	Manuel Santos (UA)
Gabriel Monteiro (IST)	Margarida Casal (UM, CBMA)
Guilherme Matos Ferreira (UA)g)	M. Leonor Nunes (IPIMAR)
Helena Gil (UCoimbra)	Miguel Prazeres (IST)
Helena Santos (ITQB/UNL)	Milton Costa (FCT/UC)
Hermínia de Lencastre (ITQB/UNL)	Nelson Lima (UM, IBB)
Ilda Sanches (FCT/UNL, CREM)	Pedro Moradas Ferreira (IBMC)
Isabel Sá Correia (IST, IBB)	Raquel Aires Barros (IST)
Ivonne Delgadillo (UA)	Rogério Tenreiro (FC/UL, BioFIG)
João Coutinho (UA)	Tim Hogg (ESB,UCP)

## Food Microbiology & Biotechnology

### P030

#### EFFECT OF TEMPERATURE ON THE BIOACTIVE PROPERTIES OF BEE POLLEN

Pascoal, A.<sup>1</sup>; Rocha, J.F.<sup>1</sup>; Choupina, A.<sup>1</sup>; Pereira, A.<sup>1</sup>; Santos, L.<sup>1</sup>; Estevinho, L.M.,<sup>1</sup>

*CIMO–Mountain Research Center. Department of Biology and Biotechnology. Agricultural College of Bragança, Polytechnic Institute of Bragança*

Bee pollen is flower pollen collected by the honey bee, *Apis mellifera*, for the purpose of feeding its larvae in the early stages of development. It is recognized to be a valuable apitherapeutic product with potential for medical, health and nutritional applications. The objective of this work was to compare the effect of different storage conditions in the bioactive compounds and biological properties of bee pollen.

The amount in phenolic compounds determined by the method of *Folin–Ciocalteu* varied from 32.64 to 48.40 mg GAE/g, for the dehydrated and refrigerated samples, respectively. Significant differences were verified among the two conservation processes. The amount in total flavonoids determined by the method *chloride of aluminum* were not verified among significant differences in the samples (6.58 mg CAE/g) and (6.99 mg CAE/g). Concerning the amount of total flavonoids determined by the method of aluminum chloride were not significant differences between the dehydrated (6.58 mg of DEA / g) and refrigerated samples (CAE 6.99 mg / g).

The antioxidant capacity of BP extracts was assessed through the scavenging effect on DPPH (2,2-diphenyl-1-picrylhydrazyl) and reducing power. The concentration-dependent antioxidative capacity was verified in DPPH and reducing power assays. Low values of EC50 on DPPH scavenging assay were obtained for fresh and dehydrated samples 0.74 and 1.16 mg/mL, respectively. For reducing power the values obtained for the refrigerated samples and dehydrated were similar. The high activity of refrigerated samples could be related with the conservation process. Regarding to phenolic compounds determined by HPLC method, were identified four (4) families, flavones being the most abundant one in the refrigerated and dehydrated samples.

We also verified that the presence of pollen differentially affected the growth of bacterium Gram-positive (*Staphylococcus xylosus*, *Staphylococcus epidermidis*), Gram-negative (*Shigella spp*, *Klebsiella pneumoniae*) and yeasts (*Candida parapsilosis*, *Candida membranifaciens*, *Candida glabrata*) under study, depending this on the microorganism and the method of BP conservation.