

M. Oliveira^{1,2}, K. Slezakova^{1,2},
M.J. Gomes³, A. Azevedo³, J.P.
Teixeira^{4,5}, C. Delerue-matos¹,
M.C. Pereira², S. Morais^{1*}

¹REQUIMTE-LAQV, Instituto Superior de Engenharia, Instituto Politécnico do Porto, R. Dr. António Bernardino de Almeida 431, Porto, Portugal.

²LAPABE, Departamento de Engenharia Química, Faculdade de Engenharia, Universidade do Porto, R. Dr. Roberto Frias, Porto, Portugal

³Escola Superior de Saúde, Instituto Politécnico de Bragança, Av. D. Afonso V, Bragança, Portugal.

⁴Instituto Nacional de Saúde Pública, Departamento de Saúde Ambiental, R. Alexandre Hercúlo 321, Porto, Portugal.

⁵Instituto de Saúde Pública, Universidade do Porto, R. das Taipas 135, Porto, Portugal.

*sbm@isep.ipp.pt

INTRODUCTION

- Firefighting occupational exposure is classified as possible carcinogen to humans by the International Agency for Research on Cancer and the US National Institute for Occupational Safety and Health [1,2].
- Tobacco smoke is a very important factor in the assessment of occupational exposure of workers, since the prolonged exposure to tobacco smoke is by itself the major cause of lung cancer [3]. The consumption of tobacco is responsible for the exposure to many smoke components including more than sixty known carcinogens, including some polycyclic aromatic hydrocarbons (PAHs) [4].
- PAHs are ubiquitous compounds formed during pyrolysis or incomplete combustion of organic matter, being well-known for their toxic, mutagenic, and carcinogenic properties to humans [5,6]. So far, the impact of tobacco smoke on firefighters' total exposure to PAHs is very limited.
- Full monitoring of firefighters' exposure to PAHs via all exposure routes should be performed through the quantification of their internal dose.
- The present study aims to perform the exposure assessment of Portuguese firefighters by six urinary PAH metabolites (OH-PAHs: 1-hydroxynaphthalene (1OHNaph), 1-hydroxyacenaphthene (1OHAc), 2-hydroxyfluorene (2OHFlu), 1-hydroxyphenanthrene (1OHPhen), 1-hydroxypyrene (1OHPy) and 3-hydroxybenzo[a]pyrene (3OHB[a]P) in smoking (S) and non-smoking (NS) firefighters.

EXPERIMENTAL

- A total of 21 healthy firefighters serving at Bragança fire station (North of Portugal) were selected through a structured questionnaire [7]. The collected information was age, weight, number of years as firefighters; participation in firefighting activities; tobacco smoking habits (including the number of cigarettes smoked per day for smokers) and on the most frequently consumed meals (boiled, roasted, and grilled) during the week before urine collection. Only firefighters that were not involved in firefighting activities within the last five days before sampling were considered. According to the information collected through the questionnaires, individuals were separated in two groups: smoking and non-smoking subjects.
- Firefighters collected a spot urine sample in sterilized 50 mL polycarbonate containers at the end of their work shift. Urine samples were frozen at -20 °C until analysis.
- Extraction and chromatographic analysis of urinary OH-PAHs were done according to the detailed description in a previous study of this research group [8]. All determination were performed in triplicate.
- The concentration of urinary creatinine was determined according to the Jaffe colorimetric method [9] and used to normalize the urinary concentrations of OH-PAHs.

Acknowledgements:

Authors are thankful to all firefighters involved in the study. This work was supported by European Union (FEDER funds through COMPETE) and National Funds (Fundação para a Ciência e Tecnologia, FCT) through projects UID/QUI/50006/2013, POCI-01-0145-FEDER-007265 and POCI-01-0145-FEDER-006939, by the FCT/MEC with national funds and co-funded by FEDER in the scope of the P2020 Partnership Agreement. Additional financial support was provided by FCT through the fellowship SFRH/BPD/105100/2014.



RESULTS

- All firefighters reported to have a diet exclusive of charbroiled and deep-fried foods within the five days before urine collection, thus making negligible the contribution of food to the overall exposure to PAHs.
- The concentrations of OH-PAHs were adjusted with the levels of creatinine for each firefighter's spot urine sample. Urinary creatinine concentrations ranged between 2,77 to 2,90 g/L for both smoking and non-smoking firefighters. No significant differences were found between the medians of these two groups ($p > 0.05$).
- 1OHNaph and/or 1OHAc were detected in 100 and 80% of the smoking and non-smoking firefighters' urines, respectively.
- 1OHPhen, 2OHFlu, and 1OHPy were detected in the urine of all firefighters.
- 3OHB[a]P was never detected.

Biometric characterization of the firefighters that participated in the study (n=21)


| Characteristic | Smoking | Non-smoking |
|--|--------------------|--------------------|
| Firefighters (n) | 6 | 15 |
| Age (mean ± SD; min – max; years) | 22 ± 1.4 (21-23) | 40 ± 5.6 (34-48) |
| Body mass index (mean ± SD; min – max) | 26 ± 2.4 (24-28) | 29 ± 4.2 (25-36) |
| Number of years as firefighters | | |
| ≤ 10 years (%) | 100 | 17 |
| 10 – 20 years (%) | 0 | 33 |
| ≥ 20 years (%) | 0 | 50 |
| Number of cigarettes smoked per day (mean ± SD; min – max) | 11 ± 1.4 (10-12) | n.a. |
| Diastolic blood pressure (mean ± SD; min – max; mmHg) | 92 ± 7.1 (87-97) | 86 ± 11 (72-106) |
| Systolic blood pressure (mean ± SD; min – max; mmHg) | 138 ± 19 (125-152) | 137 ± 11 (127-152) |
| Cardiac frequency (mean ± SD; min – max; heart beats/min.) | 79 ± 5.7 (75-83) | 72 ± 9.4 (62-84) |
| Respiratory pathologies | No | No |
| Allergies | No | No |

n.a. – not applicable



- Urinary Σ OH-PAHs were 0.738 (0.491-1.02) $\mu\text{mol/mol}$ creatinine for smoking and 0.429 (0.053-0.472) $\mu\text{mol/mol}$ creatinine for non-smoking firefighters

- Concentrations increased 72% in smoking firefighters (* $p = 0.003$, nonparametric Mann-Whitney U test)



- Smoking firefighters working at BRG fire station presented urinary levels of all individual OH-PAHs, except for 2OHFlu, 72-167% higher than in non-smokers (0.676 versus 0.394 $\mu\text{mol/mol}$ creatinine for 1OHNaph and/or 1OHAc, 0.016 versus 0.008 $\mu\text{mol/mol}$ creatinine for 1OHPhen, and 0.024 versus 0.009 $\mu\text{mol/mol}$ creatinine for 1OHPy; * $p < 0.001$, nonparametric Mann-Whitney U test).
- Spearman correlations between the levels of OH-PAHs in smoking firefighters revealed that urinary 1OHPhen concentrations were globally well correlated with the levels of other PAH metabolites ($r = 0.716$, with $p < 0.173$, except with 1OHPy).
- The relation between the urinary levels of individual and total OH-PAHs with the number of smoked cigarettes for each firefighter revealed the existence of strong and significant positive correlations in smoking firefighters ($r = 0.878$, $p < 0.021$ for Σ OH-PAHs; $r = 0.891$, $p = 0.017$ for 1OHNaph and/or 1OHAc and 2OHFlu; $r = 0.905$, $p = 0.013$ for 1OHPhen).
- Limited information concerning reference standard guidelines for urinary levels of OH-PAHs is available. The urinary levels of 1OHPy, in both smoking and non-smoking firefighters, was much lower than the guideline of 0.5 $\mu\text{mol/mol}$ creatinine recommended by the American Conference of Governmental Industrial Hygienists for occupational exposure to PAHs in smoking and non-smoking workers [10].

CONCLUSIONS

- This study proved that smoking firefighters presented significantly higher levels of urinary OH-PAHs comparatively with the control group, revealing the higher predisposition of these firefighters to develop the potential health effects associated with the exposure to PAHs.
- Regarding the incidence of cancer in firefighters, some studies reported an elevated risk of developing several types of cancers, including urothelial, skin, lung, kidney and testicular cancers [11-13] throughout their professional career; however some findings remain contradictory. Thus more studies concerning both occupational and environmental exposure of firefighters as well as toxicological and epidemiological studies are needed to better estimate the overall and cumulative risks and the potential health effects of firefighters occupational exposure.

References

- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. International Agency for Research on Cancer, 98 (2010) 405-569.
- U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention. National Institute for Occupational Safety and Health, DHHS (NIOSH) 2005-149 (2007) 1-383.
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. International Agency for Research on Cancer, 83 (2004) 1-1492.
- IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, 38 (1986) 1-432.
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, International Agency for Research on Cancer, 92 (2010) 1-853.
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, International Agency for Research on Cancer, 82, (2002) 1-556.
- WHO World Health Survey B – Individual Questionnaire. World Health Organization, Evidence and Information Policy (2002) accessed at www.who.int/healthinfo/survey/whslongindividuala.pdf (15/10/2016).
- Oliveira, M., Slezakova, K., Alves, M.J., Fernandes, A., Teixeira, J.P., Delerue-Matos, C., et al. Int. J. Hyg. Environ. Health (2016) <http://dx.doi.org/10.1016/j.ijheh.2016.07.011>.
- Kanagasabapathy, A.S. and Kumari, S. World Health Organization, Regional Office for South-East Asia, New Delhi (2000) pp. 25-28.
- ACGIH American Conference of Governmental Industrial Hygienists. Cincinnati, Ohio, USA (2010) accessed at <http://www.nsc.org/facultyportal/Documents/fih-6e-appendix-b.pdf> (
- Daniels, R.D., Kubale, T.L., Yiin, J.H., Dahm, M.M., Hales, T.R., Baris, D., et al. Occup. Environ. Med. 0, (2013) 1-10.
- LeMasters, G.K., Genaidy, A.M., Succop, P., Daddens, J., Sobehi, T., Barriera-Viruet, H., et al. J. Occup. Environ. Med. 48 (2006) 189-202.
- Golka, K., and Weistenhöfer, W. J. Tox. Env. Health B 11 (2008) 32-44.