

# HONEY AND HONEY BEES OF GUINEA-BISSAU



Pinto, M.A.<sup>1</sup>; Batista, V.<sup>1</sup>; Alves, D.<sup>2</sup>; Vilas-Boas, M.<sup>1</sup>



<sup>1</sup>Mountain Research Centre (CIMO), Polytechnic Institute of Bragança, Campus de Sta. Apolónia, Apartado 1172, 5301-855 Bragança, Portugal, [mvilasboas@ipb.pt](mailto:mvilasboas@ipb.pt)

<sup>2</sup>Apilegre, Mercado Municipal, 6050-340 Nisa, Portugal

Download this poster at [http://esa.ipb.pt/pdf/Gb\\_Q\\_f.pdf](http://esa.ipb.pt/pdf/Gb_Q_f.pdf)


## Beekeeping in Guinea – Bissau




Beekeeping is an ancient activity in Guinea-Bissau. The ancestral interaction with bees stands on “honey hunting” of natural colonies or use of traditional hives hanged on trees. These hives are perfect shelters for swarms but the colony is destroyed every year after honey harvesting. Bees are therefore kept as wild as ever with little, if any, interference from man.



## Sampling



Fifteen colonies from 7 different localities were examined for morphometry and mtDNA. Six honey samples were collected from beekeepers using Kenyan top-bar or traditional hives and analyzed for color, humidity, conductivity, free acidity, diastase activity, HMF, total phenols and main sugars using IHC methods.



## Honey Bee morphometry





The morphometric analysis showed a bee smaller than the Europeans but with similar leg and wing size. Regarding color, each worker displayed a very distinct yellow spot on thorax and a black spot at the fourth ring, which appeared either isolated from the black strip or linked, looking like a “T shape”. Those two features mark the difference towards other honey bees.

## Honey Bee Genetics

The mitochondrial DNA analysis was performed using the *Dral* test (Garnery et al. 1993), which consists in the amplification of the tRNA<sub>Leu</sub>-cox2 intergenic region followed by digestion with the restriction enzyme *Dral*.

Restriction map and fragment sizes of the tRNA<sub>Leu</sub>-cox2 intergenic region



Three different haplotypes (A1, A4, A8) were detected, all of African ancestry. As found by others for sub-Saharan Africa (Franck et al. 2001), A1 was the most common followed by A8 and A4.

## Honey quality

Harvesting Method	Color	Humidity max. 21	Conductivity max. 800	Free acidity max. 50	Fruct.+ gluc. min. 60	Diastase min. 8	HMF max. 40	Total Phenols
Honey from traditional harvesting	Dark amber	16.2%	701 $\mu\text{S}\cdot\text{cm}^{-1}$	33.5 meq. $\cdot\text{kg}^{-1}$	72%	41.4 Schade. $\cdot\text{g}^{-1}$	19.8 mg $\text{kg}^{-1}$	589 mg GAE. $\cdot\text{kg}^{-1}$
Honey from modern harvesting	Dark amber	15.3	302	12.0	60	27.3	10.8	466
	Light amber	20.4%	1015 $\mu\text{S}\cdot\text{cm}^{-1}$	32.0 meq. $\cdot\text{kg}^{-1}$	77%	22.0 Schade. $\cdot\text{g}^{-1}$	98.8 mg $\text{kg}^{-1}$	1054 mg GAE. $\cdot\text{kg}^{-1}$
Honey from modern harvesting	Amber	20.2	762	23.5	68	21.5	39.6	963
	Amber	20.1	622	17.0	60	21.0	7.7	805

References: Franck P, L Garnery, A Loiseau, BP Oldroyd, HR Hepburn, M Solignac, JM Cornuet. 2001. Genetic diversity of the honey bee in Africa: microsatellite and mitochondrial data. *Heredity* 86:420-430. Garnery L, M Solignac, G Celebrano, J M Cornuet. 1993. A simple test using restricted PCR-amplified mitochondrial DNA to study the genetic structure of *Apis mellifera* L. *Experientia* 49:1016-1021.