

Cite this: *Food Funct.*, 2016, 7, 2204

Chemical characterization and bioactive properties of *Geranium molle* L.: from the plant to the most active extract and its phytochemicals

V. C. Graça,^{a,b} Lillian Barros,^b Ricardo C. Calheta,^b Maria Inês Dias,^b Ana Maria Carvalho,^b Celestino Santos-Buelga,^c Isabel C. F. R. Ferreira^{*b} and P. F. Santos^{*d}

After a period of indifference, in which synthetic compounds were favored, there is an increasing interest in the study of the biological properties of plants and the active principles responsible for their therapeutic properties. *Geranium molle* L. has been used in the Portuguese folk medicine for the treatment of various ailments including cancer but, unlike many of the species from the *Geranium* genus, its phytochemical characterization and biological activity are virtually unexplored. In this study a *G. molle* sample from Trás-os-Montes, north-eastern Portugal, was chemically characterized regarding nutritional value, free sugars, organic acids, fatty acids and tocopherols, and several aqueous (decoction, infusion) and organic (*n*-hexane, dichloromethane, ethyl acetate, acetone, methanol) extracts of the plant were assessed for their bioactive properties. The antioxidant activity was evaluated by means of the free radicals scavenging activity, reducing power and inhibition of lipid peroxidation. The cytotoxicity of the different extracts was assessed *in vitro* against several human cancer cell lines (breast, lung, cervical and hepatocellular carcinomas) and, additionally, their hepatotoxicity was evaluated using a porcine liver primary cell culture. *G. molle* was shown to be rich in carbohydrates and proteins, providing tocopherols and essential fatty acids. Amongst the various extracts, the acetone extract was found to have the highest content of phenolic compounds (mainly ellagitannins, but also some flavone and flavonol glycosides) as well as the highest antioxidant and cytotoxic activities. To the best of our knowledge, this is the first report on the chemical composition and bioactive properties of *G. molle*.

Received 1st December 2015,

Accepted 2nd April 2016

DOI: 10.1039/c5fo01479d

www.rsc.org/foodfunction

1. Introduction

The use of medicinal plants to improve health was highly valued in ancient civilizations. Until the mid-nineteenth century, plants were the main therapeutic agents used by humans.^{1,2} After a period of indifference, in which synthetic compounds were favored, in recent years increasing interest of researchers has been observed in the study of the biological activity of plants and the active principles responsible for their therapeutic properties.^{3,4}

Geranium molle L., commonly known as Dove's-foot Crane's-bill or Dovesfoot Geranium, is an annual or biennial herb that belongs to the Geraniaceae family. This plant is native of Europe, northern Africa, and western Asia. It was introduced in southern Africa, the Americas, eastern Asia, Australia, and New Zealand.^{5,6} *Geranium molle* L. grows spontaneously in cultivated and waste places, open habitats, dunes, and dry grassland or roadsides, between 0 and 1500 m in altitude.^{7,8}

Unlike other species of the *Geranium* genus, namely the closely related *Geranium robertianum* L., there seems to be an almost complete absence of references to the use of *Geranium molle* in folk medicine. Exception is made to an ethnopharmacological study carried out by Neves *et al.*⁹ in some regions of Trás-os-Montes (north east Portugal), where the flowering aerial parts and roots of the wild plant are traditionally used to prepare decoctions and infusions for stomach acidity and stomach ache, gingivitis, eye inflammation and cuts, uterus inflammation and cancer treatment.

The phytochemistry of the *Geranium* genus is reasonably well-known and clearly dominated by phenolic constituents,¹⁰

^aChemistry Dep. and Centre for Research and Technology of Agro-Environmental and Biological Sciences (CITAB) - Vila Real, University of Trás-os-Montes and Alto Douro, 5001-801 Vila Real, Portugal

^bMountain Research Centre (CIMO), ESA, Polytechnic Institute of Bragança, Campus de Santa Apolónia, 1172, 5301-855 Bragança, Portugal. E-mail: iferreira@ipb.pt; Fax: +351-273-325405; Tel: +351-273-303219

^cGIP-USAL, Faculty of Pharmacy, University of Salamanca, Campus Miguel de Unamuno, 37007 Salamanca, Spain

^dChemistry Dep. and Chemistry Center - Vila Real (CQVR), University of Trás-os-Montes and Alto Douro, 5001-801 Vila Real, Portugal. E-mail: psantos@utad.pt; Fax: +351-259-350480; Tel: +351-259-350276

the most studied classes of compounds being tannins, flavonoids and phenolic acids. The phenolic compounds, especially the flavonoids, have been cited as the main biologically active components among those found in *Geranium* species.¹¹ This class of compounds from *Geranium* spp. were reported to exhibit antiviral, antitumor, antithrombotic, hepatoprotective, anti-inflammatory, antiallergic, antiproliferative, anticancer and immune stimulant effects.^{11–13}

Although the *Geranium* genus phytochemistry is relatively well-known, reports on the chemical composition and biological properties of *Geranium molle* L. cannot be found in the literature. Therefore, in the present study, a wild sample of *Geranium molle* L. was analysed for its nutritional composition (proteins, fat, carbohydrates and ash) and chemically characterized regarding hydrophilic (sugars, organic acids) and lipophilic (fatty acids and tocopherols) molecules. An infusion and a decoction (common forms of consumption) and different organic extracts were evaluated for their bioactive properties, namely the antioxidant and antitumor properties, being the most active extract characterized in terms of phenolic compounds.

2. Materials and methods

2.1. Standards and reagents

Acetonitrile (99.9%), *n*-hexane (97%) and ethyl acetate (99.8%) were of HPLC grade from Fisher Scientific (Lisbon, Portugal). The fatty acid methyl ester (FAME) reference standard mixture 37 (standard 47885-U) was purchased from Sigma (St Louis, MO, USA), as also, L-ascorbic acid, Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid), organic acids, sugar standards, acetic acid, formic acid, ellipticine, sulphorhodamine B (SRB), trypan blue, trichloroacetic acid (TCA) and Tris. Phenolic compound standards were purchased from Extrasynthèse (Genay, France). Tocol (50 mg mL⁻¹) and individual tocopherols were purchased from Matreya (Pleasant Gap, PA, USA). 2,2-Diphenyl-1-picrylhydrazyl (DPPH) was obtained from Alfa Aesar (Ward Hill, MA, USA). Foetal bovine serum (FBS), L-glutamine, Hank's balanced salt solution (HBSS), trypsin-EDTA (ethylenediaminetetraacetic acid), penicillin/streptomycin solution (100 U mL⁻¹ and 100 mg mL⁻¹, respectively), RPMI-1640 and DMEM media were from Hyclone (Logan, Utah, USA). Water was treated in a Milli-Q water purification system (TGI Pure Water Systems, Greenville, SC, USA).

2.2. Plant material

Geranium molle L. specimens in blossom (including thin roots, basal leaves, ascending stems, upper hairy leaves and flowers) were collected in Serra da Nogueira, Bragança, north-eastern Portugal, in March 2015, and subsequently, all dirt and dried parts were cleaned out. The amount of plant material collected was around 780 g. Voucher specimens are deposited at the herbarium of the Escola Superior Agrária de Bragança (BRESA). The botanical identification was confirmed by the

agronomist Dra. Ana Maria Carvalho of the School of Agriculture, Polytechnic Institute of Bragança (Trás-os-Montes, Portugal). The sample was lyophilized, reduced to a fine dried powder (~20 mesh), mixed to obtain a homogeneous sample and stored in a refrigerator at -20 °C, protected from light.

2.3. Chemical characterization

2.3.1. Macronutrient composition of the crude plant material. The sample was analysed for its nutritional chemical composition (proteins, fat, carbohydrates and ash) through standard procedures.¹⁴ The crude protein content (N × 6.25) of the sample was estimated by the macro-Kjeldahl method. The crude fat was determined by extracting a known weight of powdered sample with petroleum ether, using a Soxhlet apparatus. The ash content was determined by incineration at 600 ± 15 °C. Total carbohydrates were calculated by difference. Energy was calculated according to the following equation: energy (kcal) = 4 × (g protein + g carbohydrate) + 9 × (g fat).

2.3.2. Hydrophilic compounds

Free sugars. Free sugars were determined *via* high performance liquid chromatography coupled to a refraction index detector (HPLC-RI), after an extraction procedure previously described by the authors¹⁵ using melezitose as internal standard (IS). The equipment consisted of an integrated system with a pump (Knauer, Smartline system 1000, Berlin, Germany), a degasser system (Smart line manager 5000) and an auto-sampler (AS-2057 Jasco, Easton, MD, USA) coupled to a refraction index (RI) detector (Knauer Smartline 2300). The chromatographic separation was achieved with a Eurospher 100-5 NH₂ column (5 µm, 4.6 × 250 mm, Knauer) operating at 30 °C (7971 R Grace oven). The mobile phase was acetonitrile/deionized water, 70:30 (v/v), at a flow rate of 1 mL min⁻¹. Sugar identification was performed by comparing the relative retention times of sample peaks with standards. Data were analyzed using Clarity 2.4 Software (DataApex, Podohradská, Czech Republic). Quantification was based on the RI signal response of each standard, using the IS (melezitose) method and by using the calibration curves obtained for the commercial standards of each compound. The results were expressed in g per 100 g of dry weight.

Organic acids. Organic acids were determined following a procedure previously optimized and described by the authors.¹⁶ The analysis was performed using a Shimadzu 20A series ultra-fast liquid chromatograph (UFLC) (Shimadzu Corporation, Kyoto, Japan). Separation was achieved on a SphereClone (Phenomenex, Torrance, CA, USA) reverse phase C₁₈ column (5 µm, 4.6 × 250 mm) thermostatted at 35 °C. The elution was performed with 3.6 mM sulphuric acid using a flow rate of 0.8 mL min⁻¹. Detection was carried out with a diode array detector (DAD), using 215 nm and 245 nm (for ascorbic acid) as the preferred wavelengths. The organic acids found were quantified by comparison of the area of their peaks, recorded at 215 or 245 nm, with the calibration curves obtained for commercial standards of each compound. The results were expressed in g per 100 g of dry weight.

2.3.3. Lipophilic compounds

Fatty acids. Fatty acids were determined after transesterification according to the procedure previously described by the authors.¹⁵ The fatty acid profile was analyzed with a DANI 1000 gas chromatograph (GC) equipped with a split/splitless injector, a flame ionization detector (FID) and a Macherey-Nagel (Düren, Germany) column (50% cyanopropyl-methyl-50% phenylmethylpolysiloxane, 30 m × 0.32 mm i.d. × 0.25 μm d_f). The oven temperature program was as follows: the initial temperature of the column was 50 °C, held for 2 min, then a 30 °C per min ramp to 125 °C, 5 °C per min ramp to 160 °C, 20 °C per min ramp to 180 °C, 3 °C per min ramp to 200 °C, 20 °C per min ramp to 220 °C and held for 15 min. The carrier gas (hydrogen) flow-rate was 4.0 mL min⁻¹ (0.61 bar), measured at 50 °C. Split injection (1:40) was carried out at 250 °C. Fatty acid identification was performed by comparing the relative retention times of the sample's FAME peaks with standards. The results were recorded and processed using Clarity 4.0.1.7 Software (DataApex, Podohradska, Czech Republic) and expressed as a relative percentage of each fatty acid.

Tocopherols. Tocopherols were determined following a procedure previously described by the authors.¹⁵ The analysis was performed by HPLC (equipment described in Section 2.3.2, sub section Free sugars), using a fluorescence detector (FP-2020; Jasco, Easton, MD, USA) programmed for excitation at 290 nm and emission at 330 nm. The chromatographic separation was achieved with a Polyamide II (YMC Waters, Milford, MA, USA) normal-phase column (5 μm, 4.6 mm × 250 mm), operating at 35 °C. The mobile phase used was a mixture of *n*-hexane and ethyl acetate (70:30, v/v) at a flow rate of 1 mL min⁻¹. The compounds were identified by chromatographic comparison with authentic standards. Quantification was based on the fluorescence signal response of each standard, using the IS (tocol) method, and the calibration curves obtained for commercial standards of each compound. The results were expressed in mg per 100 g of dry weight.

2.4. Preparation of organic and aqueous extracts

The organic (hexane, dichloromethane, ethyl acetate, acetone and methanol) and aqueous (obtained by infusion and decoction) extracts were prepared from the lyophilized plant.

For the preparation of the organic extracts a sample (100 g) was extracted with 500 mL of *n*-hexane and the mixture was stirred vigorously at room temperature, 150 rpm, for 48 h, and then filtered under reduced pressure successively through a Whatman No. 541 paper and a sintered glass funnel. The solid residue was extracted with an additional 500 mL of *n*-hexane under the same conditions. The combined extracts were evaporated to dryness at 40 °C under reduced pressure. The resulting residue was further extracted sequentially with dichloromethane, ethyl acetate, acetone and finally methanol, according to the procedure described above.

The infusions were prepared by adding the sample (1 g) to 100 mL of boiling distilled water, left to stand at room temperature for 5 min, and then filtered under reduced pressure

successively through a Whatman No. 541 paper and a sintered glass funnel. The obtained infusions were frozen and lyophilized.

The decoctions were also prepared by adding the sample (1 g) to 100 mL of distilled water and boiled for 5 min. The mixture was left to stand for 5 min at room temperature and then filtered under reduced pressure successively through a Whatman No. 541 paper and a sintered glass funnel. The obtained decoctions were frozen and lyophilized.

2.5. Bioactive compounds in the extracts

For total phenolics determination, an aliquot of the different extracts (1 mL, 78–625 μg mL⁻¹) were mixed separately, with Folin-Ciocalteu reagent (5 mL, previously diluted with water 1:10 v/v) and sodium carbonate (75 g L⁻¹, 4 mL). The tubes were vortexed for 15 s and allowed to stand for 30 min at 40 °C for color development. Absorbance was then measured at 765 nm.¹⁷ Gallic acid was used to calculate the standard curve (0.1–1 mM) and the results were expressed as mg of gallic acid equivalents (GAE) per g of extract.

For total flavonoids determination, an aliquot of the different extracts (0.5 mL, 313–625 μg mL⁻¹) were mixed separately, with distilled water (2 mL) and subsequently with NaNO₂ solution (5%, 0.15 mL). After 6 min, AlCl₃ solution (10%, 0.15 mL) was added and allowed to stand further 6 min, thereafter, NaOH solution (4%, 2 mL) was added to the mixture. Immediately, distilled water was added to bring the final volume to 5 mL. Then the mixture was mixed properly and allowed to stand for 15 min. The intensity of the pink color was measured at 510 nm.¹⁸ Catechin was used to calculate the standard curve (0.3–1 mM) and the results were expressed as mg of catechin equivalents (CE) per g of extract.

2.6. Evaluation of bioactive properties of the extracts

2.6.1. Antioxidant activity assays. The organic extracts were redissolved in methanol (final concentration 10 mg mL⁻¹). The aqueous extracts were redissolved in water (final concentration 10 mg mL⁻¹). The final solutions obtained were further diluted to different concentrations to be subjected to distinct evaluation assays of the antioxidant activity.

DPPH radical-scavenging activity was evaluated by using an ELX800 microplate reader (Bio-Tek Instruments, Inc.; Winooski, VT, USA), and calculated as a percentage of DPPH discoloration using the formula: $[(A_{\text{DPPH}} - A_{\text{S}})/A_{\text{DPPH}}] \times 100$, where A_{S} and A_{DPPH} are, respectively, the absorbance of the sample solution and that of the DPPH solution at 515 nm. Reducing power was evaluated by the capacity to convert Fe³⁺ into Fe²⁺, measuring the absorbance at 690 nm in the microplate reader mentioned above. Inhibition of β-carotene bleaching was evaluated through the β-carotene/linoleate assay; the neutralization of linoleate free radicals avoids β-carotene bleaching, which is measured by the formula: $(\beta\text{-carotene absorbance after 2 h of assay}/\text{initial absorbance}) \times 100$. Lipid peroxidation inhibition in porcine (*Sus scrofa*) brain homogenates was evaluated by the decrease in thiobarbituric acid reactive substances (TBARS); the color intensity of the

malondialdehyde–thiobarbituric acid (MDA–TBA) was measured by its absorbance at 532 nm; the inhibition ratio (%) was calculated using the following formula: $[(A - B)/A] \times 100\%$, where A and B are the absorbance of the control and the sample solution, respectively.^{16,19}

The results were expressed in EC₅₀ values (sample concentration providing 50% of antioxidant activity or 0.5 of absorbance in the reducing power assay). Trolox was used as positive control.

2.6.2. Cytotoxicity in human tumor cell lines. The aqueous and organic extracts were redissolved in water and ethanol 20%, respectively, in order to obtain a final concentration of 8 mg mL⁻¹. The final solution was further diluted to different concentrations (400 to 1.5 μg mL⁻¹) to be subjected to *in vitro* cytotoxicity evaluation. Four human tumor cell lines were used: MCF-7 (breast adenocarcinoma), NCI-H460 (non-small cell lung cancer), HeLa (cervical carcinoma) and HepG2 (hepatocellular carcinoma). The cells were routinely maintained as adherent cell cultures in RPMI-1640 medium containing 10% heat-inactivated FBS (MCF-7 and NCI-H460) and 2 mM glutamine or in DMEM supplemented with 10% FBS, 2 mM glutamine, 100 U per mL penicillin and 100 mg per mL streptomycin (HeLa and HepG2 cells), at 37 °C, in a humidified air incubator containing 5% CO₂. Each cell line was plated at an appropriate density (7.5 × 10³ cells per well for MCF-7 and NCI-H460 or 1.0 × 10⁴ cells per well for HeLa and HepG2) in 96-well plates and allowed to attach for 24 h. The cells were then treated for 48 h with the different diluted sample solutions. Following this incubation period, the adherent cells were fixed by adding cold 10% TCA (100 μL) and incubated for 60 min at 4 °C. Plates were then washed with deionized water and dried; SRB solution (0.1% in 1% acetic acid, 100 μL) was then added to each plate-well and incubated for 30 min at room temperature. Unbound SRB was removed by washing with 1% acetic acid. Plates were air-dried, the bound SRB was solubilised with 10 mM Tris (200 μL, pH 7.4) and the absorbance was measured at 540 nm in the microplate reader mentioned above.¹⁶ The results were expressed in GI₅₀ values (sample concentration that inhibited 50% of the net cell growth). Ellipticine was used as positive control.

2.6.3. Hepatotoxicity in non-tumor cells. A cell culture was prepared from a freshly harvested porcine liver obtained from a local slaughter house, and it was designed as PLP2. Briefly, the liver tissues were rinsed in Hank's balanced salt solution containing 100 U per mL penicillin, 100 μg per mL streptomycin and divided into 1 × 1 mm³ explants. Some of these explants were placed in 25 cm² tissue flasks in DMEM medium supplemented with 10% fetal bovine serum, 2 mM nonessential amino acids and 100 U per mL penicillin, 100 mg per mL streptomycin and incubated at 37 °C under a humidified atmosphere containing 5% CO₂. The medium was changed every two days. Cultivation of the cells was continued with direct monitoring every two to three days using a phase contrast microscope. Before confluence was reached, cells were subcultured and plated in 96-well plates at a density of 1.0 × 10⁴ cells per well, and cultivated in DMEM medium with 10%

FBS, 100 U per mL penicillin and 100 μg per mL streptomycin.^{15,16} Cells were treated for 48 h with the different diluted sample solutions and the same procedure described in the previous section for the SRB assay was followed. The results were expressed in GI₅₀ values (sample concentration that inhibited 50% of the net cell growth). Ellipticine was used as positive control.

2.7. Phenolic composition of the acetone extract

The acetone extract was redissolved in water/methanol 80:20 (v/v) (final concentration 5 mg mL⁻¹). Phenolic compounds were determined by HPLC (Hewlett-Packard 1100 chromatographer, Agilent Technologies, Santa Clara, CA, USA) as previously described by the authors.^{16,19} Double online detection was carried out with a DAD using 280 nm and 370 nm as preferred wavelengths and with a mass spectrometer (MS) connected to the HPLC system *via* the DAD cell outlet. The phenolic compounds were identified by comparing their retention time, UV-vis and mass spectra with those obtained from standard compounds, when available. Otherwise, peaks were tentatively identified comparing the obtained information with the available data reported in the literature. For quantitative analysis, a calibration curve for each available phenolic standard was constructed based on the UV signal. For the identified phenolic compounds for which a commercial standard was not available, the quantification was performed through the calibration curve of other compound from the same phenolic group. The results were expressed in mg per g of extract.

2.8. Statistical analysis

For all the experiments, three samples were analyzed and all the assays were carried out in triplicate. The results are expressed as mean values ± standard deviation (SD). The differences between the different samples were analyzed using one-way analysis of variance (ANOVA) followed by Tukey's honestly significant difference *post hoc* test with $\alpha = 0.05$, coupled with Welch's statistic. This treatment was carried out using the SPSS v. 22.0 program.

3. Results and discussion

3.1. Chemical characterization of *Geranium molle* L.

The results of the nutritional characterization of *G. molle*, namely, macronutrients, sugars, organic acids and fatty acids, are presented in Table 1. Carbohydrates were the most abundant macronutrients found in the studied sample, followed by proteins, fat and ash. This plant showed 72.2% of moisture and the energetic contribution was ~436 kcal per 100 g dw.

Fructose, glucose and sucrose were the free sugars detected in this sample. The total sugar content was ~9 g per 100 g dw, glucose being present in a much larger amount (~6 g per 100 g dw) than the two other sugars.

A total of six different organic acids were detected in the plant, namely, oxalic, quinic, malic, ascorbic, citric and fumaric acids (Table 1). Malic acid was the most abundant

Table 1 Chemical characterization of *Geranium molle* L. in terms of macronutrients, and hydrophilic and lipophilic compounds

	Quantity		Quantity
<i>Macronutrients</i>			
Moisture (g per 100 g fw)	72.2 ± 0.3	C14:1	0.14 ± 0.01
Fat (g per 100 g dw)	15.5 ± 0.5	C15:0	0.95 ± 0.01
Proteins (g per 100 g dw)	20.9 ± 0.4	C16:0	24.43 ± 0.01
Ash (g per 100 g dw)	10.5 ± 0.1	C16:1	0.69 ± 0.01
Carbohydrates (g per 100 g dw)	53.1 ± 0.2	C17:0	1.21 ± 0.01
Energy (kcal per 100 g dw)	436 ± 2	C18:0	17.2 ± 0.2
		C18:1n9	26.1 ± 0.1
		C18:2n6	4.58 ± 0.07
		C18:3n3	5.56 ± 0.01
<i>Hydrophilic compounds</i>			
Fructose	1.62 ± 0.01	C20:0	0.72 ± 0.02
Glucose	6.32 ± 0.01	C20:1	0.08 ± 0.01
Sucrose	0.99 ± 0.01	C20:2	0.034 ± 0.002
Sum of sugars (g per 100 g dw)	8.93 ± 0.02	C20:3n6	0.08 ± 0.01
Oxalic acid	0.71 ± 0.01	C20:4n6	0.39 ± 0.01
Quinic acid	2.6 ± 0.3	C20:3n3 + C21:0	0.18 ± 0.01
Malic acid	2.821 ± 0.002	C20:5n3	0.20 ± 0.01
Ascorbic acid	0.362 ± 0.001	C22:0	0.47 ± 0.01
Citric acid	1.09 ± 0.05	C22:1n9	0.038 ± 0.001
Fumaric acid	0.0063 ± 0.0001	C22:6n3	0.22 ± 0.01
Sum of organic acids (g per 100 g dw)	7.6 ± 0.3	C24:0	0.30 ± 0.01
		C24:1	0.043 ± 0.001
<i>Lipophilic compounds</i>			
C6:0	2.11 ± 0.01	SFA (%)	61.64 ± 0.18
C8:0	1.39 ± 0.01	MUFA (%)	27.12 ± 0.09
C10:0	3.35 ± 0.01	PUFA (%)	11.24 ± 0.08
C11:0	0.022 ± 0.001	α-Tocopherol	18.9 ± 0.6
C12:0	1.98 ± 0.01	β-Tocopherol	0.33 ± 0.01
C13:0	0.063 ± 0.004	γ-Tocopherol	0.76 ± 0.01
C14:0	7.39 ± 0.01	Sum of tocopherols (mg per 100 g dw)	19.99 ± 0.55

(C6:0), caproic acid; (C8:0), caprylic acid; (C10:0), capric acid; (C11:0), undecylic acid; (C12:0), lauric acid; (C13:0), tridecanoic acid; (C14:0), myristic acid; (C14:1), myristoleic acid; (C15:0), pentadecanoic acid; (C16:0), palmitic acid; (C16:1), palmitoleic acid; (C17:0), heptadecanoic acid; (C18:0), stearic acid; (C18:1n9c+t), oleic acid; (C18:2n6c), linoleic acid; (C18:3n3), α-linolenic acid; (C20:0), arachidic acid; (C20:1), *cis*-11-eicosenoic acid; (C20:2), *cis*-11,14-Eicosadienoic acid; (C20:3n6), eicosatrienoic acid; (C20:4n6), arachidonic acid; (C20:3n3 + C21:0), *cis*-11,14,17-eicosatrienoic acid and heneicosanoic acid; (C20:5n3), eicosapentaenoic acid; (C22:0), behenic acid; (C22:1n9), Erucic acid; (C22:6n3), dicosahexanoic acid; (C24:0), lignoceric acid; (C24:1), nervonic acid; SFA – saturated fatty acids; MUFA – monounsaturated fatty acids; PUFA – polyunsaturated fatty acids; fw – fresh weight; dw – dry weight.

(~2.8 g per 100 g dw), followed by quinic acid (~2.6 g per 100 g dw).

Twenty-eight fatty acids (FA) were determined in *G. molle* (Table 1). Highest percentages were found for oleic (C18:1n9), palmitic (C16:0) and stearic acids (C18:0). Saturated fatty acids (SFA) predominated over monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA).

Regarding tocopherols, α-tocopherol (18.9 mg per 100 g dw) was the most abundant isoform present in *G. molle*. β-Tocopherol and γ-tocopherol were present but in very small amounts. δ-Tocopherol was not detected in this plant.

3.2. Bioactive compounds in different *Geranium molle* L. extracts

Concentrations of total polyphenols and total flavonoids, as determined by photometrical methods in different *G. molle* extracts are presented in Table 2. The acetone extract displayed the highest content of both total polyphenols (497 mg GAE per g extract) and total flavonoids (112 mg CE per g extract). Acetone has been reported as a good solvent for the extraction of phenolic compounds and flavonoids.^{20,21} The infusion and the decoction presented similar concentrations of total flavo-

noids but somewhat different content of total polyphenols, the extract resulting from the infusion being richer in this group of compounds. The dichloromethane and the *n*-hexane extracts showed very low concentrations of both total polyphenols and total flavonoids.

3.3. Bioactive properties of different *Geranium molle* L. extracts

The *in vitro* antioxidant and cytotoxic properties of different extracts of *G. molle* were evaluated, and the results are presented in Table 3. The antioxidant activity was determined by free radical (DPPH) scavenging activity, reducing power, inhibition of lipid peroxidation in brain cell homogenates, and TBARS assays. The cytotoxicity was tested against human tumor cell lines (breast, lung, cervical and hepatocellular carcinomas) and the hepatotoxicity was evaluated using a porcine liver primary cell culture.

In general, all the extracts revealed antioxidant potential. The acetone extract displayed the highest antioxidant activity in all the assays. This was probably related to the higher content of total phenols and total flavonoids found in this extract, compared to the other prepared extracts (Table 2). The

Table 2 Bioactive compounds in different *Geranium molle* L. extracts

Extracts	Aqueous extracts		Organic extracts				
	Infusion	Decoction	<i>n</i> -Hexane	Dichloromethane	Ethyl acetate	Acetone	Methanol
Total Polyphenols (mg GAE per g extract)	79 ± 1c	63 ± 1d	13 ± 1e	6.15 ± 0.03f	216 ± 2b	497 ± 8a	76 ± 5c
Total Flavonoids (mg CE per g extract)	27 ± 1d	25.1 ± 0.2d	4.5 ± 0.4e	2.1 ± 0.02e	74 ± 6b	112 ± 1a	53 ± 3c

GAE – gallic acid equivalents; CE – catechin equivalents. In each row different letters mean significant differences ($p < 0.05$).

Table 3 Bioactive properties of different *Geranium molle* L. extracts

	Aqueous extracts		Organic extracts				
	Infusion	Decoction	<i>n</i> -Hexane	Dichloromethane	Ethyl acetate	Acetone	Methanol
Antioxidant activity (EC ₅₀ , µg mL ⁻¹)							
DPPH scavenging activity	324 ± 9b	248 ± 4c	1816 ± 126a	>10 000	128 ± 5d	18.9 ± 0.5e	135 ± 3d
Reducing power	141 ± 1c	170 ± 6b	266 ± 5a	265 ± 1a	51 ± 1e	20.3 ± 0.2f	105 ± 4d
β-Carotene bleaching inhibition	197 ± 8e	249 ± 9b	226 ± 4c	253 ± 11b	212 ± 5d	61 ± 3f	274 ± 6a
TBARS inhibition	54 ± 3d	144 ± 7a	98 ± 4c	130 ± 6b	34 ± 2e	6.5 ± 0.2f	38 ± 2e
Antitumor activity (GI ₅₀ values, µg mL ⁻¹)							
MCF-7 (breast carcinoma)	225 ± 15b	187 ± 9c	>400	370 ± 20a	215 ± 14b	85 ± 13d	229 ± 17b
NCI-H460 (non-small lung cancer)	190 ± 16bc	172 ± 3c	>400	256 ± 17a	200 ± 14b	63 ± 4d	206 ± 8b
HeLa (cervical carcinoma)	226 ± 19b	267 ± 18a	211 ± 18b	234 ± 21b	232 ± 15b	56 ± 5c	204 ± 15b
HepG2 (hepatocellular carcinoma)	241 ± 14a	170 ± 13c	>400	>400	204 ± 16b	50 ± 4d	260 ± 13a
Hepatotoxicity (GI ₅₀ value, µg mL ⁻¹)							
PLP2	>400	>400	>400	>400	338 ± 12a	191 ± 15b	332 ± 32a

The antioxidant activity was expressed as EC₅₀ values, which means that higher values correspond to lower reducing power or antioxidant potential. EC₅₀: extract concentration corresponding to 50% of antioxidant activity or 0.5 of absorbance in reducing power assay. Trolox EC₅₀ values: 41 µg mL⁻¹ (reducing power), 42 µg mL⁻¹ (DPPH scavenging activity), 18 µg mL⁻¹ (β-carotene bleaching inhibition) and 23 µg mL⁻¹ (TBARS inhibition). GI₅₀ values correspond to the sample concentration achieving 50% of growth inhibition in human tumour cell lines or in liver primary culture PLP2. Ellipticine GI₅₀ values: 1.21 µg mL⁻¹ (MCF-7), 1.03 µg mL⁻¹ (NCI-H460), 0.91 µg mL⁻¹ (HeLa), 1.10 µg mL⁻¹ (HepG2) and 2.29 µg mL⁻¹ (PLP2). In each row different letters mean significant differences ($p < 0.05$).

antioxidant properties of many plants are closely related to the presence of phenolic compounds, which constitute the most abundant secondary metabolites of plants.²²

Regarding the cytotoxic properties, almost all of the extracts showed activity. The aqueous extract obtained by infusion, and methanol and ethyl acetate extracts showed similar cytotoxic effects against MCF-7, NCI-H460 and HeLa cells. The dichloromethane extract was ineffective against the HepG2 cell line even at the maximum concentration assayed (400 µg mL⁻¹). The *n*-hexane extract showed activity only against HeLa cells. The acetone extract displayed the highest cytotoxic effect, being significantly more potent than the remaining extracts (GI₅₀ values approximately four times lower) against all cell lines. Unfortunately, this extract also presented the highest toxicity against normal primary cells from the porcine liver (PLP2) (GI₅₀ ~ 190 µg mL⁻¹). However, the concentration required to reach 50% of growth inhibition of PLP2 is about 3–4 times higher than the concentration required to achieve 50% of growth inhibition of the human tumor cell lines tested. Therefore, a dosage of 85 µg mL⁻¹ of the *G. molle* acetone extract would be recommended to guarantee 50% of growth inhibition of the human tumor cell lines tested, without presenting toxicity effects for non-tumor cells.

The aqueous, dichloromethane and *n*-hexane extracts did not show hepatotoxicity against PLP2 cells up to the maximal tested concentration (GI₅₀ > 400 µg mL⁻¹). The methanol and ethyl acetate extracts presented similar hepatotoxicity against this cell line (GI₅₀ ~ 330 µg mL⁻¹).

Trolox and ellipticine were used as positive controls in the antioxidant and cytotoxic activity assays, respectively. However, as these are individual compounds, they should not be considered as standards and comparison with the results obtained for the extracts/oral preparations should be avoided since an eventual synergistic effect of the mixtures cannot be precluded.

3.4. Analysis of phenolic compounds in the acetone extract

Amongst the several *G. molle* extracts prepared, the acetone extract was found to exhibit the highest amount of total phenolic compounds, as well as the highest antioxidant and cytotoxic activity, so that its detailed phenolic composition was also analysed. The HPLC phenolic profile of that extract recorded at 280 and 370 nm is shown in Fig. 1. The peak characteristics and tentative identities are presented in Table 4. Sixteen phenolic compounds were detected, five of which were ellagitannins, one phenolic acid and ten flavonoids.




Fig. 1 Phenolic profile of *Geranium molle* L. acetone extract recorded at 370 nm (A) and 280 nm (B).

Table 4 Retention time (R_t), wavelengths of maximum absorption in the visible region (λ_{\max}), mass spectral data, identification and quantification of phenolic compounds of the acetone extract of *Geranium molle* L.

Peak	R_t (min)	λ_{\max} (nm)	Molecular ion $[M - H]^-$ (m/z)	MS^2 (m/z)	Tentative identification	Quantification ($mg\ g^{-1}$)
1	4.7	278	633	463(5), 421(3), 301(82), 275(49), 169(8)	Galloyl-HHDP-glucose	4.8 ± 0.1
2	6.1	276	785	633(9), 615(10), 483(15), 301(64), 275(30), 169(7)	Digalloyl-HHDP-glucose	61 ± 4
3	7.7	278	289	245(40), 203(30), 187(40), 161(30), 137(20)	(+)-Catechin	16 ± 1
4	7.9	328	353	191(100), 179(47), 173(39), 161(20), 135(25)	5- <i>O</i> -Caffeoylquinic acid	7 ± 1
5	8.4	276	785	633(6), 615(8), 483(20), 301(77), 275(28), 169(8)	Digalloyl-HHDP-glucose	66 ± 5
6	12.5	276	783	765(77), 721(5), 481(5), 301(13), 275(17)	Bis-HHDP-glucose	43 ± 3
7	14.1	278	783	765(89), 721(3), 481(3), 301(17), 275(15)	Bis-HHDP-glucose	58 ± 4
8	15.5	358	755	301(100)	Quercetin- <i>O</i> -dideoxyhexosyl-hexoside	0.18 ± 0.01
9	16.0	350	447	429(15), 357(90), 327(80), 297(50), 285(21)	Luteolin-6- <i>C</i> -glucoside	4.7 ± 0.3
10	17.0	358	609	301(100)	Quercetin-3- <i>O</i> -rutinoside	0.37 ± 0.03
11	18.9	340	563	443(17), 413(33), 311(67), 293(33)	Apigenin 2''- <i>O</i> -pentosyl-6- <i>C</i> -hexoside	0.69 ± 0.04
12	19.7	336	431	413(5), 341(40), 311(100), 283(53)	Apigenin-6- <i>C</i> -glucoside	1.53 ± 0.10
13	20.1	358	463	301(100)	Quercetin-3- <i>O</i> -glucoside	0.60 ± 0.02
14	21.1	354	593	285(100)	Kaempferol-3- <i>O</i> -rutinoside	0.89 ± 0.01
15	23.1	356	433	301(100)	Quercetin- <i>O</i> -pentoside	0.20 ± 0.02
16	24.1	348	447	285(100)	Kaempferol-3- <i>O</i> -glucoside	0.36 ± 0.02
					Total ellagitannins	234 ± 16
					Total hydroxycinnamoyl esters	7 ± 1
					Total flavonoids	25 ± 2
					Total phenolic compounds	265 ± 19

(+)-Catechin (compound 3), 5-*O*-caffeoylquinic acid (compound 4), luteolin-6-*C*-glucoside (compound 9), quercetin-3-*O*-rutinoside (compound 10), apigenin-6-*C*-glucoside (compound 12), quercetin-3-*O*-glucoside (compound 13), kaempferol-3-*O*-rutinoside (compound 14) and kaempferol-3-*O*-glucoside (compound 16) were positively identified according to their retention, mass and UV-vis characteristics by comparison with commercial standards.

Compounds 1, 2 and 5–7 were identified as hydrolysable tannins (ellagitannin derivatives) and assigned as different hexahydroxydiphenyl (HHDP)-glucose esters based on their pseudomolecular ions and fragmentation patterns. Thus, the

mass characteristics of compound 1 ($[M - H]^-$ at m/z 633) pointed to a galloyl-HHDP-glucose isomer, whereas those of compounds 2 and 5 ($[M - H]^-$ at m/z 785) were consistent with digalloyl-HHDP-glucose isomers, and compounds 6 and 7 with bis-HHDP-glucose isomers.^{23,24} Similar compounds have been described in other *Geranium* species;^{10,25–31} actually HHDP esters are considered as the main hydrolysable tannins and majority phenolic compounds in most species of *Geranium*.^{10,25,27,32,33} However, to the best of our knowledge, they have never been reported in *G. molle*.

The remaining compounds would correspond to flavonoids. Compounds 8 ($[M - H]^-$ at m/z 755) and 15 ($[M - H]^-$ at m/z

447) were associated to quercetin glycosides based on their absorption spectra and the production of an ion fragment at m/z 301. Their molecular weights pointed to a quercetin-*O*-pentoside (peak 15) and a quercetin derivative bearing two deoxyhexosyl and one hexosyl residues (peak 8). In the latter derivative, the fact that only one MS^2 fragment was released corresponding to the aglycone suggests that the three sugars constituted a trisaccharide. Flavonol derivatives have been previously described in other *Geranium* sp. pl., namely different quercetin and kaempferol aglycones and glycoside derivatives.^{34–40}

Finally compound 11 ($[M - H]^-$ at m/z 563) was tentatively assigned as apigenin 2''-*O*-pentosyl-6-*C*-hexoside according to its pseudomolecular ion and fragmentation pattern. A compound with the same characteristics had been previously found by our group in *Arenaria montana*⁴¹ and identified based on the fragmentation patterns described by Ferreres *et al.*⁴² and Ferreres *et al.*⁴³ for *O,C*-glycosyl flavones. As far as we know this type of compound is reported herein for the first time in *Geranium* species, although some flavone *O*-glycosides and aglycones, *i.e.*, luteolin and apigenin, have been reported in *Geranium* sp. pl.^{35,38–40}

Digalloyl-HHDP-glucoside (compound 5) was the most abundant ellagitannin present, while luteolin-6-*C*-glucoside (compound 9) was the best represented flavonoid. Chlorogenic acid (*i.e.*, 5-*O*-caffeoylquinic acid) was the only hydroxycinnamoyl derivative found in the analysed sample.

4. Conclusions

Geranium molle L. is rich in carbohydrates and proteins, providing tocopherols and essential fatty acids. In general, the various aqueous and organic extracts showed antioxidant activity and cytotoxicity against the different human tumour cell lines tested. The acetone extract presented the highest antioxidant potential in the different assays, which is most probably related to its higher content of polyphenols and flavonoids compared with the other prepared extracts. The acetone extract also displayed the highest cytotoxic effect, being significantly more potent than the remaining extracts against all cell lines (GI_{50} values approximately four times lower). Although this extract also presented the highest toxicity against porcine liver primary cells (PLP2), the GI_{50} value for PLP2 was about 3–4 times higher than those for the tumor cell lines tested. The decoction and the infusion of the plant, which are the common forms of folk consumption, did not show hepatotoxicity against PLP2 cells up to the maximal tested concentration (400 $\mu\text{g mL}^{-1}$), but presented GI_{50} for the tumor cells 3–4 times higher than the acetone extract. The phenolic profile of the acetone extract was determined by HPLC-MS and shown to be constituted mainly by ellagitannins, as well as some flavone and flavonol glycosides. All in all, the obtained results support the folk medicinal use of *G. molle*, and its interest as a source of phytochemicals with bioactive properties to be explored in the medicine and food industries.

Acknowledgements

This work is supported by national funds by FCT – Portuguese Foundation for Science and Technology, under the project UID/AGR/04033/2013 and PEst-OE/AGR/UI0690/2014. V. C. Graça, M. I. Dias, R. C. Calhella and L. Barros thank FCT for SFRH/BD/52542/2014, SFRH/BD/84485/2012, SFRH/BPD/68344/2010 and SFRH/BPD/107855/2015 grants, respectively.

References

- 1 J. Camejo-Rodrigues, L. Ascensão, A. M. Bonet and J. Vallès, *J. Ethnopharmacol.*, 2003, **89**, 199–209.
- 2 J. L. Longe, *The Gale Encyclopedia of Alternative Medicine*, Thomson Gale, Farmington Hills, 2nd edn, 2005.
- 3 N. C. C. Silva and A. Fernandes-Júnior, *J. Venom. Anim. Toxins incl. Trop. Dis.*, 2010, **16**, 402–413.
- 4 L. Rubió, M.-J. Motilva and M.-P. Romero, *Crit. Rev. Food Sci. Nutr.*, 2013, **53**, 943–953.
- 5 C. Aedo, J. J. Aldasoro and C. Navarro, *Ann. Missouri Bot. Gard.*, 1998, **85**, 594–630.
- 6 C. Aedo, F. M. Garmendia and F. Pando, *Anales Jard. Bot. Madrid*, 1998, **56**, 211–252.
- 7 S. Pignatti, *Flora d'Italia – Edagricole*, 1982, vol. II, p. 10.
- 8 C. Aedo, *Anales Jard. Bot. Madrid*, 2000, **58**, 39–82.
- 9 J. M. Neves, C. Matos, C. Moutinho, G. Queiroz and L. R. Gomes, *J. Ethnopharmacol.*, 2009, **124**, 270–283.
- 10 J. B. Harborne and C. A. Williams, in *Geranium and Pelargonium – The genera Geranium and Pelargonium*, ed. M. Lis-Balchin, Taylor & Francis, New York, 2002, vol. 4, pp. 20–29.
- 11 E. Neagu, G. P. Roman, G. L. Radu and G. Nechifor, *Rom. Biotechnol. Lett.*, 2010, **15**, 5042–5048.
- 12 A. Saija, M. Scalase, M. Lanza, D. Marzulla, F. Bonina and F. Castelli, *Free Radical Biol. Med.*, 1995, **19**, 481–486.
- 13 J.-U. Shim, P.-S. Oha and K.-T. Limb, *J. Ethnopharmacol.*, 2009, **126**, 90–95.
- 14 AOAC, *Official methods of analysis*, Association of Official Analytical Chemists Gaithersburg, Maryland, 18th edn, 2005.
- 15 L. Barros, E. Pereira, R. C. Calhella, M. Dueñas, A. M. Carvalho, C. Santos-Buelga and I. C. F. R. Ferreira, *J. Funct. Foods*, 2013, **5**, 1732–1740.
- 16 M. I. Dias, L. Barros, M. Dueñas, E. Pereira, A. M. Carvalho, R. C. Alves, M. B. P. P. Oliveira, C. Santos-Buelga and I. C. F. R. Ferreira, *Food Chem.*, 2013, **141**, 4152–4160.
- 17 V. L. Singleton, R. Orthofer and R. M. Lamuela-Raventos, *Methods Enzymol.*, 1999, **299**, 152–178.
- 18 J. Zhishen, T. Mengcheng and W. Jianming, *Food Chem.*, 1999, **64**, 555–559.
- 19 C. L. Roriz, L. Barros, A. M. Carvalho, C. Santos-Buelga and I. C. F. R. Ferreira, *Food Res. Int.*, 2014, **62**, 684–693.
- 20 L. Barros, M. Dueñas, I. C. F. R. Ferreira, P. Baptista and C. Santos-Buelga, *Food Chem. Toxicol.*, 2009, **47**, 1076–1079.

- 21 J. Azmir, I. S. M. Zaidul, M. M. Rahman, K. M. Sharif, A. Mohamed, F. Sahena, M. H. A. Jahurul, K. Ghafoor, N. A. N. Norulaini and A. K. M. Omar, *J. Food Eng.*, 2013, **117**, 426–436.
- 22 J. Dai and R. J. Mumper, *Molecules*, 2010, **15**, 7313–7352.
- 23 P. Mena, L. Calani, C. Dall'Asta, G. Galaverna, C. García-Viguera, R. Bruni, A. Crozier and D. Del Rio, *Molecules*, 2012, **17**, 14821–14840.
- 24 J. Regueiro, C. Sánchez-González, A. Vallverdú-Queralt, J. Simal-Gándara, R. Lamuela-Raventós and M. Izquierdo-Pulido, *Food Chem.*, 2014, **152**, 340–348.
- 25 E. C. Bate-Smith, *Phytochemistry*, 1972, **11**, 1755–1757.
- 26 T. Okuda, K. Mori, K. Seno and T. Hatano, *J. Chromatogr.*, 1979, **171**, 313–320.
- 27 E. C. Bate-Smith, *Phytochemistry*, 1981, **20**, 211–216.
- 28 T. Okuda, T. Yoshida and T. Hatano, *Phytochemistry*, 2000, **55**, 513–529.
- 29 T. Okuda, *Phytochemistry*, 2005, **66**, 2012–2031.
- 30 A. Tuominen, *Phytochemistry*, 2013, **95**, 408–420.
- 31 A. Tuominen, E. Toivonen, P. Mutikainen and J.-P. Salminen, *Phytochemistry*, 2013, **95**, 394–407.
- 32 T. Okuda, K. Mori and T. Hatano, *Phytochemistry*, 1980, **19**, 547–551.
- 33 Q.-Y. Wu, Y. Zhou, X. Jin, Y. Guan, M. Xu and L.-F. Liu, *Int. J. Mol. Sci.*, 2011, **12**, 8740–8749.
- 34 T. Kartnig and J. Bucar-Stachel, *Planta Med.*, 1991, **57**, 292–293.
- 35 S. Ivancheva and A. Petrova, *Biochem. Syst. Ecol.*, 2000, **28**, 255–260.
- 36 K. B. Kobakhidze and M. D. Alaniya, *Chem. Nat. Compd.*, 2004, **40**, 89–90.
- 37 C. S. Fodorea, L. Vlase, S. Suciuc, M. Tamas and S. E. Leucuta, *Rev. Med. Chir. Soc. Med. Nat. Iasi*, 2005, **109**, 174–178.
- 38 G. Paun, E. Neagu, S. C. Litescu, P. Rotinberg and G. L. Radu, *J. Serb. Chem. Soc.*, 2012, **77**, 1191–1203.
- 39 E. Neagu, G. Paun, D. Constantin and G. L. Radu, *Arab. J. Chem.*, 2013, DOI: 10.1016/j.arabjc.2013.09.028.
- 40 G. Paun, S. C. Litescu, E. Neagu, A. Tache and G. L. Radu, *J. Enzyme Inhib. Med. Chem.*, 2014, **29**, 28–34.
- 41 E. Pereira, L. Barros, R. C. Calhelha, M. Dueñas, A. M. Carvalho, C. Santos-Buelga and I. C. F. R. Ferreira, *Food Funct.*, 2014, **5**, 1848–1855.
- 42 F. Ferreres, A. Gil-Izquierdo, P. B. Andrade, P. Valentão and F. A. Tomás-Barberán, *J. Chromatogr. A*, 2007, **1161**, 214–223.
- 43 F. Ferreres, A. Gil-Izquierdo, J. Vinholes, C. Grosso, P. Valentão and P. B. Andrade, *Rapid Commun. Mass Spectrom.*, 2011, **25**, 700–712.