

KINEMATICAL CHARACTERISATION OF A BASIC HEAD-OUT AQUATIC EXERCISE DURING AN INCREMENTAL PROTOCOL

INTRODUCTION

Massive research has been produced throughout the last decades in order to better understand the role of head-out aquatic exercises in populations' health (Barbosa et al, 2009). Indeed, such studies aimed to characterize the physiological acute and/or chronic response of subjects performing head-out aquatic exercises. Moreover, the comprehensive knowledge about the biomechanical (i.e. kinematical) behavior performing this aquatic activity is quite reduced.

Conducting head-out aquatic exercise sessions, instructors often use the music cadence to achieve a pre-imposed intensity of exertion. Music cadences between 130 and 150 b.min⁻¹ are suggested by technical literature for head-out aquatic exercises (Kinder and See, 1992). At least one empirical study reported that for healthy and physically active subjects instructors should choose music cadences between 136 and 158 b.min⁻¹ (Barbosa et al., 2010).

Increases in the music cadence imposed significant increases in the acute physiological adaptation (e.g., rate of perceived effort, heart rate or blood lactate) of the subjects (Hoshijima et al., 1999; Barbosa et al., 2010). It is hypothesized that increased physiological response may be explained by the fact that increasing music cadence will also increase movement velocity and frequency, decreasing the segmental range of motion. However, to our knowledge there is no study in the literature reporting the kinematical changes imposed by an incremental cadence protocol at head-out aquatic exercises.

The aim of this study was to analyze the relationships between musical cadence and kinematical characteristics of a basic head-out aquatic exercise, when immersed to the xiphoid process (i.e., breast).




Figure 1. The basic aquatic exercise studied and called "rocking horse".

METHODS

Six non-pregnant, clinically healthy and physically active young women holding a graduation degree in Sports Sciences and at least one year of experience conducting head-out aquatic classes volunteered to participate in this study (23.67 ± 0.52 y-old; 57.42 ± 4.78 kg of body mass; 1.64 ± 0.07 m of height; 22.17 ± 2.56 kg.m⁻² of body mass index).

The protocol consisted of five bouts of 16 repetitions performing the basic head-out aquatic exercise "rocking horse" (Figure 1) at the "water tempo" immersed to the xiphoid process (i.e., breast). Bouts intensity were 80 %, 90 %, 100 %, 110 % and 120 % of the cadence reported by Barbosa et al. (2010) to achieve a 4 mmol.l⁻¹ of blood lactate, representing respectively 120 b.min⁻¹, 135 b.min⁻¹, 150 b.min⁻¹, 165 b.min⁻¹ and 180 b.min⁻¹ cadences. Musical cadence was controlled electronically by a metronome (Korg, MA-30, Tokyo, Japan) connected to a sound system.

The protocol was videotaped in sagittal plane with a pair of cameras providing a dual projection from both underwater (GR-SXM25 SVHS, JVC, Yokoama, Japan) and above (GR-SX1 SVHS, JVC, Yokoama, Japan) the water surface. The images of both cameras were recorded independently. The study comprised the kinematical analysis of the full cycles (Ariel Performance Analysis System, Ariel Dynamics Inc., USA) through a VCR (Panasonic, AG 7355, Japan) at a frequency of 50 Hz. Zatsiorsky's model with an adaptation by de Leva (1996) was used with the division of the trunk in two articulated parts. To create a single image of dual projection as described previously (Vilas-Boas et al., 1997; Barbosa et al., 2005), the independent digitalization from both cameras was reconstructed with the help of a calibration object (0.675 x 0.855 m; 6 control points) and a 2D-DLT algorithm (Abdel-Aziz and Karara, 1971). For the analysis of the curve of the centre of mass's kinematics a filter with a cut-off frequency of 5 Hz was used, as suggested by Winter (1990). For the segmental kinematics a cut-off frequency of 9 Hz was used, near to the value proposed by Winter (1990). A double-pass filtering for the signal processing was used.

METHODS

It was evaluated the: (i) cycle period; (ii) 2D linear position ranges (foot, hand and centre of mass); 2D linear velocity ranges (foot, hand and centre of mass).

The normality of the distributions was assessed with the Shapiro-Wilk test. Linear regression equations models and its coefficients of determination were used to describe the relationships between musical cadence and biomechanical variables. The level of statistical significance was set at P ≤ 0.05.




Figure 2. Qualitative analysis from the centre of mass' horizontal Kinematics'.




Figure 3. Simple scatter gram from the cycle period according to the cadence imposed.




Figure 4. Overlay scatter gram from horizontal displacement and vertical displacement according to the cadence imposed.




Figure 5. Overlay scatter gram from horizontal velocity and vertical velocity according to the cadence imposed.

RESULTS AND DISCUSSION

Figure 2 presents a qualitative analysis from the centre of mass kinematics from a single subject during the second bout at 135 b.min⁻¹. Figure 3 reports the simple scatter gram from the cycle period according to the musical cadence imposed. Figure 4 and 5 presents respectively the overlay scatter gram for 2D displacement and 2D velocity according to the musical cadence imposed.

CONCLUSION

As a conclusion, expert and fit subjects seem to increase segmental velocity with increasing cadences to avoid the decrease of the segmental range of motion.

REFERENCES

- Abdel-Aziz Y, Karara, H. (1971). Direct linear transformation: from comparator coordinates into object coordinates in close range photogrammetry. Proceedings of the Symposium on close-range photogrammetry, p. 1-18, Church Falls.
- Barbosa TM, Keskinen KL, Fernandes RJ, Colação P, Lima AB, Vilas-Boas JP. (2005). Energy cost and intracyclic variation of the velocity of the centre of mass in butterfly stroke. *Eur J Appl Phys* 93:519-523.
- Barbosa TM, Marinho DA, Bragada JA, Reis VM, Silva AJ. (2009). Physiological assessment of head-out aquatic exercises in healthy subjects: a qualitative review. *J Sports Sci Med*. 8: 179-189
- Barbosa TM, Sousa V, Silva AJ, Reis V, Marinho DA, Bragada JA. (2010). Effects of music cadence in the acute physiological adaptations to head-out aquatic exercises. *J Strength Cond Res*. 24: 244-250
- de Leva P. (1996). Adjustments to Zatsiorsky-Seluyanov's segment inertia parameters. *Journal of Biomechanics* 29:1223-1230.
- Hoshijima Y, Torigoe K, Yamamoto M, Nishimura S, Abo N, Imoto M, Miyachi M, Onodera S. (1999). Effects of music rhythm on heart rate and oxygen uptake during squat exercises in water and on land. *Biomechanics and Medicine in Swimming VIII*, p. 337-339. Gummerus Printing.
- Kinder T, See J. (1992). *Aqua Aerobics - A Scientific Approach*. Dubuque: Eddie Bowers Publishing
- Vilas-Boas JP, Cunha P, Figueiras T, Ferreira M, Duarte J. (1997). Movement analysis in simultaneous swimming techniques. *Cologne Swimming Symposium*. Verlag. p.95-103. Sport Fahnemann
- Winter D (1990). *Biomechanic and Motor Control of Human Movement*. Chichester: John Wiley & Sons.